

The Eph Receptor/Ephrin System: An Emerging Player in the Invasion Game

Tessa N. Campbell and Stephen M. Robbins

Southern Alberta Cancer Research Institute,
Departments of Oncology and Biochemistry & Molecular
Biology, University of Calgary, Calgary, AB

Abstract

Eph receptor tyrosine kinases (Ephs) and their membrane-anchored ligands (ephrins) form a vital cell communication system capable of bi-directional signaling. This Eph receptor/ephrin system has classically been demonstrated to play a role in development. However, emerging evidence has revealed differential expression of Ephs and ephrins in numerous cancers. Recent studies suggest that this system influences invasive behaviour, promoting a more aggressive and metastatic phenotype. Hence, this mini-review summarizes the current understanding of the contribution of both Eph receptors and their ephrin ligands to invasiveness in cancer, as well as their use as potential therapeutic targets.

Eph Receptor/ Ephrin Signaling

Eph receptors (erythropoietin-producing human hepatocellular carcinoma) comprise the largest family of vertebrate receptor tyrosine kinases. In concert with their ephrin ligands (Eph family receptor interacting proteins), they form an essential cell-cell communication system capable of bi-directional signaling, where receptor signaling is designated "forward" and ephrin signaling is "reverse" (Fig. 1; Kullander and Klein, 2002; Heroult et al., 2006).




Fig. 1. Bi-directional signaling in the Eph receptor/ephrin communication system.

Eph receptors have been divided into an EphA subclass (8 members) and an EphB subclass (6 members) on the basis of sequence similarity and ligand affinity (Gale et al., 1996). Ephrin ligands have also been divided into

two subclasses: GPI-linked ephrin As (5 members) and transmembrane ephrin Bs (3 members). Although promiscuity has been observed, the ephrin A ligands preferentially bind to EphA receptors, while ephrin B ligands bind preferentially to EphB receptors (Heroult et al., 2006).

The Eph receptor/ephrin system has been demonstrated to play a role in numerous biological processes. Generally, Ephs and ephrins are thought to act as graded molecular tags which translate the density of their cognate partner on opposing membranes into precisely graded cellular responses, resulting in cell-contact repulsion or cell-cell adhesion. Through this mechanism, the Eph receptor/ephrin system has been shown to direct the positioning, adhesion and migration of cells and cell layers during development (Wimmer-Kleikamp and Lackmann, 2005). This system has also been implicated in immune regulation (Wu and Luo, 2005), as well as in central nervous system injury and disease (Goldshmit et al., 2006). Recently, a role for the Eph receptor/ephrin system has also emerged in cancer, especially in the area of invasive behaviour.

Eph Receptors and Ephrins in Cancer

Though the majority of research has studied the Eph/ephrin role in development, new evidence suggests strong involvement in tumourigenesis, including metastasis, angiogenesis, and invasion (Dodolet and Pasquale, 2000; Wimmer-Kleikamp and Lackmann, 2005). Upon identification of EphA1, the first member of the Eph family, breast, liver, lung and colon tumours showed overexpression, but no evidence of gene amplification nor mutation (Hirai, et al., 1987; Dodolet and Pasquale, 2000; Surawska et al., 2004). Genes for other Eph receptors and ephrins have since been recognized to be overexpressed or differentially expressed in numerous human cancers (for review, see Surawska et al., 2004). Mounting evidence documents a strong correlation between the expression levels of many Eph/ephrin family members and increased invasiveness of a number of aggressive tumours, including breast cancer, colon carcinoma, malignant melanoma, kidney carcinoma, ovarian cancer, neuroblastomas, and prostate cancer (Wimmer-Kleikamp and Lackmann, 2005). It has also been suggested that elevated Eph/ephrin expression levels may be diagnostic for specific invasive, metastatic tumours and reduced patient survival rates (Saito et al., 2004; Nakada et al., 2006).

Eph Receptors and Ephrins in Invasion

For the purpose of this review, invasion is defined as "penetration of tissue barriers, such as basement membrane and interstitial stroma, by cells" (Friedl and Wolf, 2003). It occurs during normal cell morphogenesis, wound healing, and in malignant cells. Numerous Eph receptors and ephrins have been noted to influence or correlate with invasive behaviour. Cytotrophoblast commitment to uterine invasion, for example, was noted

to be accompanied by upregulation of ephrin B1 and downregulation of EphB4 (Red-Horse et. al., 2005). Additionally, ephrin A5 expression was found to increase invasive behaviour, anchorage-independent growth, and morphological transformation of murine fibroblasts (Campbell et. al., 2006). The majority of research on Ephs and ephrins in invasion, however, has focused on their role in cancer (Table 1).

With respect to the EphA/ephrin A subsystem, Saito et. al. (2004) noted that expression level of EphA2 had a statistically significant relationship with liver metastasis, lymphatic vessel invasion, and clinical stage. Alford et. al. (2007) found that transglutaminase-cross-linked ephrin A1 and A5 bound to EphA receptors and promoted invasion and migration of HeLa cells. Moreover, Iida et. al. (2005) reported that the poor prognosis of patients with AFP-producing hepatocellular carcinoma was partially caused by ephrin A1 expression, which induced expression of genes related to tumour cell growth, angiogenesis, invasion, and metastasis.

With respect to involvement of the EphB/ephrin B subsystem in invasion, a greater depth of investigation has occurred, resulting in both observational and mechanistic insight. Nakada et. al. (2004) demonstrated that EphB2-transfected glioma cells showed increased invasive behaviour both *in vitro* and in an *ex vivo* rat brain slice. Furthermore, glioma invasion was promoted by activation of EphB2 or inhibited by blocking EphB2. This invasive behaviour was revealed to be controlled through an EphB2/R-Ras signaling pathway (Nakada

et. al., 2005).

EphB4 was shown to confer survival and invasive properties in head and neck squamous cell carcinoma (Masood et. al., 2006). Moreover, Alam et. al. (2007) reported that histoscores and mRNA levels of EphB4 and ephrin B2 significantly increased with clinical stages, dedifferentiation, and myometrial invasion in uterine endometrial cancer. Similarly, Takai et. al. (2004) noted that ephrin B2 and EphB4 expression were significantly associated with the presence of clinical stage, histological grade, and invasion to >1/2 myometrium in endometrial carcinomas. Currently, controversy surrounds the role of EphB4 in invasiveness and potential prognosis in breast cancer, since this receptor has been reported to act both as a tumour suppressor and a tumour promoter (Noren et. al., 2006; Noren and Pasquale, 2007).

Aside from the observations described in endometrial carcinomas, ephrin B2 mRNA expression was further reported to be elevated in higher stage neoplasms and in patients with high-risk AMES in malignant thyroid cancer (Kebebew et. al., 2005). Ephrin B2 was also visualized in the invasive front of malignant melanoma (Meyer et. al., 2005).

In the case of the correlation of ephrin B1 and B3 expression with invasive properties in cancer, mechanistic insight has also been provided. For ephrin B1, tyrosine phosphorylation of this ligand promoted invasion of gastric scirrhous carcinoma cells both *in vitro* and *in vivo* (Tanaka et. al., 2007a). Through the

Table 1. Invasive involvement of Eph receptors and ephrin ligands in cancer.

Ligand/ Receptor	Cell Line(s)/ Tissue(s)	Observation(s)	Reference(s)
ephrin A1	hepatocellular carcinoma; HeLa cells	poor prognosis of patients with AFP-producing hepatocellular carcinoma is partially caused by ephrin A1 expression, which induces expression of genes related to tumour cell growth, angiogenesis, invasion, and metastasis; transglutaminase-cross-linked ephrin A1 and A5 bind to A-type Eph receptors and promote invasion and migration of HeLa cells	Iida et. al., 2005; Alford et. al., 2007
ephrin A5	HeLa cells	transglutaminase-cross-linked ephrin A1 and A5 bind to A-type Eph receptors and promote invasion and migration of HeLa cells	Alford et. al., 2007
ephrin B1	gastric scirrhous carcinoma; pancreas cancer cell lines	tyrosine phosphorylation of ephrin B1 promotes invasion of cancer cells <i>in vitro</i> and <i>in vivo</i> ; expression of ephrin B1 promotes the invasion of cancer cells <i>in vivo</i> , which requires the C-terminus of ephrin B1	Tanaka et. al., 2007a; Tanaka et. al., 2007b
ephrin B2	endometrial carcinoma; malignant thyroid neoplasms; malignant melanoma; uterine endometrial cancer	ephrin B2 and EphB4 expression are significantly associated with the presence of clinical stage, histological grade, and invasion to >1/2 myometrium; ephrin B2 mRNA expression is elevated in higher TNM stage neoplasms and in patients with high-risk AMES; prominent expression of ephrin B2 is noted in the invasive front of malignant melanoma; histoscores and mRNA levels of ephrinB2 and EphB4 significantly increase with clinical stages, dedifferentiation and myometrial invasion	Takai et. al., 2004; Kebebew et. al., 2005; Meyer et. al., 2005; Alam et. al., 2007
ephrin B3	invasive glioma cells; gliomas	ephrin B3 is expressed in invasive glioma cells; ephrin B3 expression, phosphorylation, and signaling through Rac1 are critically important to glioma invasion	Hoelzinger et. al., 2005; Nakada et. al., 2006
EphA2	colorectal cancer	expression level of EphA2 has a statistically significant relationship with liver metastasis, lymphatic vessel invasion and clinical stage	Saito et. al., 2004
EphB2	gliomas; glioma cells; colorectal cancer	U251 cells stably transfected with EphB2 show more pronounced invasive growth in an <i>ex vivo</i> rat brain slice and glioma cell invasion is promoted by activation of EphB2 or inhibited by blocking EphB2; EphB2/R-Ras signaling regulates glioma cell adhesion, growth, and invasion; reduced expression of EphB2 parallels invasion and metastasis in colorectal tumours	Nakada et. al., 2004; Nakada et. al., 2005; Guo et. al., 2006
EphB4	endometrial carcinoma; breast cancer; head and neck squamous cell carcinoma; uterine endometrial cancer	ephrin B2 and EphB4 expression are significantly associated with the presence of clinical stage, histological grade, and invasion to >1/2 myometrium; EphB4 activates an antioncogenic Abl-Crk pathway that inhibits breast cancer cell viability, proliferation, motility and invasion; histoscores and mRNA levels of ephrinB2 and EphB4 significantly increase with clinical stages, dedifferentiation and myometrial invasion; expression of EphB4 in HNSCC tumor cells confers survival and invasive properties	Takai et. al., 2004; Noren et. al., 2006; Masood et. al., 2006; Alam et. al., 2007
EphB6	breast cancer	downregulation of EphB6 correlates with the most invasive aggressive cell line and may serve as a prognostic indicator in breast cancer	Fox and Kandpal, 2004

use of multiple cell lines, including pancreatic cancer cells, it was demonstrated that the C-terminus of ephrin B1 was responsible for conferring the increased invasive properties through activation of matrix metalloproteinase-8 secretion (Tanaka et al., 2007b). For ephrin B3, increased expression was noted specifically in invasive glioma cells (Hoelzinger et al., 2005), and this elevated ephrin B3 expression accompanied by phosphorylation leading to signaling through Rac1 was identified as critically important to glioma invasion (Nakada et al., 2006).

It has been proposed that Ephs operate during tumour progression through uncontrolled re-emergence of their developmental cell guidance capacity, promoting tumour metastasis, invasion and neo-angiogenesis (Wimmer-Kleikamp and Lackmann, 2005). Interestingly, some studies have noted a down-regulation of Eph receptor expression corresponding to an invasive phenotype in certain cancers. For example, Fox and Kandpal (2004) reported that decreased expression of EphB6, which was down-regulated in the most aggressive breast cancer cell line, was particularly important in invasiveness. Additionally, Guo et al. (2006) observed that reduced expression of EphB2 paralleled invasion and metastasis in colorectal tumours. Moreover, Battle et al. (2005) also noted that loss of EphB2 in colorectal cancer correlated with degree of malignancy. Thus, researchers have chosen to focus on individual Ephs and ephrins in specific types of cancer when designing potential therapies, instead of working from a general model.

Therapeutic Implications

Based upon the observed differential expression of Eph receptors and their ephrin ligands in a wide variety of cancers, it is not surprising that components of this system have been flagged as potential therapeutic targets. One approach has been antibody targeting of Ephs identified as consistently upregulated in specific cancers, thus offering selective targeting of tumour cells while minimizing harm to healthy tissues. This method has been used to target EphA2 in breast cancer cells (Carles-Kinch et al., 2002) and EphB2 in colorectal cancer (Mao et al., 2004).

Another approach is to use antagonistic peptides that display selective binding. This method has been used in the case of EphB2 (Chrencik et al., 2007) and EphA2 (Koolpe et al., 2002). Recently, Yamaguchi et al. (2007) engineered a vaccine using dendritic cells pulsed with EphA2-derived peptides that resulted in long-term antitumour immunity against a rechallenge with MC38 tumour cells in a murine colon cancer model.

Function-blocking soluble Eph receptors have also been used to interfere with tumour growth and properties. As with the peptide approach, this method has been demonstrated for both the EphA and EphB subclasses (Brantley et al., 2002; Martiny-Baron et al., 2004; Kertesz et al., 2006; Yang et al., 2006). A recent encouraging result was reported by Dobrzanski et al. (2004) who found that administration of EphA2/Fc profoundly inhibited the growth of primary tumours and

the development of peritoneal, lymphatic, and hepatic metastases in a murine pancreatic ductal adenocarcinoma model. Interestingly, this approach was also employed from a ligand perspective by introducing soluble ephrin A1-Fc to suppress growth of EphA2-expressing gastric cancer cells (Nakamura et al., 2005).

Eph/ephrin-influenced cytoskeletal signal transduction pathways dealing with cell motility, tumour invasiveness and metastatic potential have also been investigated as therapeutic targets. Such flagged components have included FAK, CRK, Rho, paxillin, Cbl, and Grb4 (Surawska et al., 2004). Further understanding of the Eph/ephrin system will bring forth more candidate targets.

Summary

Due to the emergence of the Eph receptor/ephrin system as a contributor to cellular invasive behaviour, it is likely that this system will remain a high priority in oncology for the development of new therapies and treatments. For example, a currently underutilized approach is the use of RNA interference, the sequence-specific gene silencing induced by double-stranded RNA (Campbell and Choy, 2005). Numerous studies have utilized RNA interference to explore the effect of an Eph or ephrin knock-down *in vitro*, but have yet to engineer a successful *in vivo* method for the treatment of cancer.

Further studies in the manipulation of the Eph receptor/ephrin system will aid in engineering cancer therapies and in elucidating the complexity and additional functional implications of the system.

Acknowledgements

This work was supported by an operating grant from CIHR to S.M.R. S.M.R. currently holds a Canada Research Chair in Cancer Biology and is an Alberta Heritage Foundation for Medical Research (AHFMR) Scientist. T.N.C. was supported by AHFMR and Alberta Cancer Foundation Postdoctoral Fellowships.

References

- Alam, S.M., Fujimoto, J., Jahan, I., Sato, E., and Tamaya, T. (2007). Overexpression of ephrinB2 and EphB4 in tumor advancement of uterine endometrial cancers. *Ann. Oncol.* *18*, 485-490.
- Alford, S.C., Bazowski, J., Lorimer, H., Elowe, S., and Howard, P.L. (2007). Tissue transglutaminase clusters soluble A-type ephrins into functionally active high molecular weight oligomers. *Exp. Cell. Res.* *313*, 4170-4179.
- Battle, E., Bacani, J., Begthel, H., Jonkeer, S., Gregorieff, A., van de Born, M., Malats, N., Sancho, E., Boon, E., Pawson, T., Gallinger, S., Pals, S., and Clevers, H. (2005). EphB receptor activity suppresses colorectal cancer progression. *Nature* *435*, 1126-1130.
- Brantley, D.M., Cheng, N., Thompson, E.J., Lin, Q., Brekken, R.A., Thorpe, P.E., Muraoka, R.S., Cerretti, D.P., Pozzi, A., Jackson, D., Lin, C., and Chen, J.

- (2002). Soluble Eph A receptors inhibit tumor angiogenesis and progression in vivo. *Oncogene* 21, 7011-7026.
- Campbell, T.N., Attwell, S., Arcellana-Panlilio, M., and Robbins, S.M. (2006). Ephrin A5 expression promotes invasion and transformation of murine fibroblasts. *Biochem. Biophys. Res. Comm.* 350, 623-628.
- Campbell, T.N. and Choy, F.Y.M. (2005). RNA interference: past, present and future. *Curr. Issues Mol. Biol.* 7, 1-6.
- Carles-Kinch, K., Kilpatrick, K.E., Stewart, J.C., and Kinch, M.S. (2002). Antibody targeting of the EphA2 tyrosine kinase inhibits malignant cell behaviour. *Cancer Res.* 62, 2840-2847.
- Chrencik, J.E., Brooun, A., Recht, M.I., Nicola, G., Davis, L.K., Abagyan, R., Widmer, H., Pasquale, E.B., and Kuhn, P. (2007). Three-dimensional structure of the EphB2 receptor in complex with an antagonistic peptide reveals a novel mode of inhibition. *J Biol. Chem.* 282, 36505-36513.
- Dobrzanski, P., Hunter, K., Jones-Bolin, S., Chang, H., Robinson, C., Pritchard, S., Zhao, H., and Ruggeri, B. (2004). Antiangiogenic and antitumor efficacy of EphA2 receptor antagonist. *Cancer Res.* 64, 910-919.
- Dodelet, V.C. and Pasquale, E.B. (2000). Eph receptors and ephrin ligands: embryogenesis to tumorigenesis. *Oncogene* 19, 5614-5619.
- Fox, B.P. and Kandpal, R.P. (2004). Invasiveness of breast carcinoma cells and transcript profile: Eph receptors and ephrin ligands as molecular markers of potential diagnostic and prognostic application. *Biochem. Biophys. Res. Comm.* 318, 882-892.
- Friedl, P. and Wolf, K. (2003). Tumour-cell invasion and migration: diversity and escape mechanisms. *Nat. Rev. Cancer* 3, 362-374.
- Gale, N.W., Holland, S.J., Valenzuela, D.M., Flenniken, A., Pan, L., Ryan, T.E., Henkemeyer, M., Strebhardt, K., Hirai, H., Wilkinson, D.G., Pawson, T., Davis, S., and Yancopoulos, G.D. (1996). Eph receptors and ligands comprise two major specificity subclasses and are reciprocally compartmentalized during embryogenesis. *Neuron* 17, 9-19.
- Goldshmit, Y., McLenachan, S., and Turnley, A. (2006). Roles of Eph receptors and ephrins in the normal and damaged adult CNS. *Brain Res. Rev.* 52, 327-345.
- Guo, D.L., Zhang, J., Yuen, S.T., Tsui, W.Y., Chan, A.S., Ho, C., Ji, J., Leung, S.Y., and Chen, X. (2006). Reduced expression of EphB2 that parallels invasion and metastasis in colorectal tumours. *Carcinogenesis* 27, 454-464.
- Heroult, M., Schaffner, F., and Augustin, H.G. (2006). Eph receptor and ephrin ligand-mediated interactions during angiogenesis and tumour progression. *Exper. Cell Res.* 312, 642-650.
- Hirai, H., Maru, Y., Hagiwara, K., Nishida, J., and Takaku, F. (1987). A novel putative tyrosine kinase receptor encoded by the eph gene. *Science* 238, 1717-1720.
- Hoelzinger, D.B., Mariani, L., Weis, J., Woyke, T., Berens, T.J., McDonough, W.S., Sloan, A., Coons, S.W., and Berens, M.E. (2005). Gene expression profile of glioblastoma multiforme invasive phenotype points to new therapeutic targets. *Neoplasia* 7, 7-16.
- Iida, H., Honda, M., Kawai, H.F., Yamashita, T., Shiota, Y., Wang, B.C., Miao, H., and Kaneko, S. (2005). Ephrin-A1 expression contributes to the malignant characteristics of {alpha}-fetoprotein producing hepatocellular carcinoma. *Gut* 54, 843-851.
- Kebebew, E., Peng, M., Reiff, E., Duh, Q.Y., Clark, O.H., and McMillan, A. (2005). Diagnostic and prognostic value of angiogenesis-modulating genes in malignant thyroid neoplasms. *Surgery* 138, 1102-1109.
- Kertesz, N., Krasnoperov, V., Reddy, R., Leshanski, L., Kumar, S.R., Zozulya, S., and Gill, P.S. (2006). The soluble extracellular domain of EphB4 (sEphB4) antagonizes EphB4-EphrinB2 interaction, modulates angiogenesis, and inhibits tumor growth. *Blood* 107, 2330-2338.
- Koolpe, M., Dail, M., and Pasquale, E.B. (2002). An ephrin mimetic peptide that selectively targets the EphA2 receptor. *J. Biol. Chem.* 277, 46974-46979.
- Kullander, K. and Klein, R., (2002). Mechanisms and functions of Eph and ephrin signaling. *Nat. Rev. Mol. Cell. Biol.* 3, 475-486.
- Mao, W., Luis, E., Ross, S., Silva, J., Tan, C., Crowley, C., Chui, C., Franz, G., Senter, P., Koeppen, H., and Polakis, P. (2004). EphB2 as a therapeutic antibody drug target for the treatment of colorectal cancer. *Cancer Res.* 64, 781-788.
- Martiny-Baron, G., Korff, T., Schaffner, F., Esser, N., Eggstein, S., Marme, D., and Augustin, H.G. (2004). Inhibition of tumour growth and angiogenesis by soluble EphB4. *Neoplasia* 6, 248-257.
- Meyer, S., Hafner, C., Guba, M., Flegel, S., Geissler, E.K., Becker, B., Koehl, G.E., Orso, E., Landthaler, M., and Vogt, T. (2005). Ephrin-B2 overexpression enhances integrin-mediated ECM-attachment and migration of B16 melanoma cells. *Int. J. Oncol.* 27, 1197-1206.
- Masood, R., Kumar, S.R., Sinha, U.K., Crowe, D.L., Krasnoperov, V., Reddy, R.K., Zozulya, S., Singh, J., Xia, G., Broek, D., Schönthal, A.H., and Gill, P.S. (2006). EphB4 provides survival advantage to squamous cell carcinoma of the head and neck. *Int. J. Cancer* 119, 1236-1248.

- Nakada, M., Drake, K.L., Niska, J.A., and Berens, M.E. (2006). Ephrin-B3 ligand promotes glioma invasion through activation of Rac1. *Cancer Res.* *66*, 8492-8500.
- Nakada, M., Niska, J.A., Miyamori, H., McDonough, W.S., Wu, J., Sato, H., and Berens, M.E. (2004). The phosphorylation of EphB2 receptor regulates migration and invasion of human glioma cells. *Cancer Res.* *64*, 3179-3185.
- Nakada, M., Niska, J.A., Tran, N.L., McDonough, W.S., and Berens, M.E. (2005). EphB2/R-Ras signaling regulates glioma cell adhesion, growth, and invasion. *Am. J. Pathol.* *167*, 565-576.
- Noren, N.K., Foos, G., Hauser, C.A., and Pasquale, E.B. (2006). The EphB4 receptor suppresses breast cancer cell tumorigenicity through an Abl-Crk pathway. *Nat. Cell. Biol.* *8*, 815-825.
- Noren, N.K. and Pasquale, E.B. (2007). Paradoxes of the EphB4 receptor in cancer. *Cancer Res.* *67*, 3994-3997.
- Red-Horse, K., Kapidzic, M., Zhou, Y., Feng, K.T., Singh, H., and Fisher, S.J. (2005). EPHB4 regulates chemokine-evoked trophoblast responses: a mechanism for incorporating the human placenta into the maternal circulation. *Development* *132*, 4097-4106.
- Saito, T., Masuda, N., Miyazaki, T., Kanoh, K., Suzuki, H., Shimura, T., Asao, T., and Kuwano, H. (2004). Expression of EphA2 and E-cadherin in colorectal cancer: correlation with cancer metastasis. *Oncol. Rep.* *11*, 605-611.
- Surawska, H., Ma, P.C., and Salgia, R. (2004). The role of ephrins and Eph receptors in cancer. *Cytokine Growth Factor Rev.* *15*, 419-433.
- Takai, N., Ueda, T., Nishida, M., Nasu, K., and Miyakawa, I. (2004). The relationship between oncogene expression and clinical outcome in endometrial carcinoma. *Curr. Cancer Drug Targets* *4*, 511-520.
- Tanaka, M., Kamata, R., Takigahira, M., Yanagihara, K., and Sakai, R. (2007a). Phosphorylation of ephrin-B1 regulates dissemination of gastric scirrhous carcinoma. *Am. J. Pathol.* *171*, 68-78.
- Tanaka, M., Sasaki, K., Kamata, R., and Sakai, R. (2007b). The C-terminus of ephrin-B1 regulates metalloproteinase secretion and invasion of cancer cells. *J. Cell. Sci.* *120*, 2179-2189.
- Wimmer-Kleikamp, S.H. and Lackmann, M. (2005). Eph-modulated cell morphology, adhesion and motility in carcinogenesis. *IUBMB Life* *57*, 421-431.
- Wu, J. and Luo, H. (2005). Recent advances on T-cell regulation by receptor tyrosine kinases. *Curr. Opin. Hematol.* *12*, 292-297.
- Yamaguchi, S., Tatsumi, T., Takehara, T., Sakamori, R., Uemura, A., Mizushima, T., Ohkawa, K., Storkus, W.J., and Hayashi, N. (2007). Immunotherapy of murine colon cancer using receptor tyrosine kinase EphA2-derived peptide-pulsed dendritic cell vaccines. *Cancer* *110*, 1469-1477.
- Yang, N.Y., Pasquale, E.B., Owen, L.B., and Ethell, I.M. (2006). The EphB4 receptor-tyrosine kinase promotes the migration of melanoma cells through Rho-mediated actin cytoskeleton reorganization. *J. Biol. Chem.* *281*, 32574-32586.

Caliciviruses

Molecular and Cellular Virology

Edited by: G.S. Hansman, J. Jiang, K.Y. Green
c. 250 pp., April 2010

ISBN: 978-1-904455-63-9 \$310 / £159

The most important research findings.
Timely and comprehensive reviews.
Discussion of past and current research.

Epstein-Barr Virus

Latency and Transformation

Edited by: Erle S. Robertson
c. 220 pp., April 2010

ISBN: 978-1-904455-62-2 \$310 / £159

Expert virologists comprehensively review this important subject from a genetic, biochemical, immunological, and cell biological perspective. Essential reading.

Anaerobic Parasitic Protozoa

Genomics and Molecular Biology

Edited by: C.G. Clark, P.J. Johnson, R.D. Adam
c. 210 pp., March 2010

ISBN: 978-1-904455-61-5 \$310 / £159

Internationally acclaimed researchers critically review the most important aspects of research on anaerobic parasitic protozoa.

Lentiviruses and Macrophages

Molecular and Cellular Interactions

Edited by: Moira Desport
c. 410 pp., March 2010

ISBN: 978-1-904455-60-8 \$310 / £159

Top lentivirus and macrophage specialists comprehensively review cutting-edge topics in the molecular and cellular biology of the lentivirus-macrophage interaction.

Microbial Population Genetics

Edited by: Jianping Xu
c. 230 pp., March 2010

ISBN: 978-1-904455-59-2 \$310 / £159

Details the major current advances in microbial population genetics and genomics.

Borrelia

Molecular Biology, Host Interaction and Pathogenesis

Edited by: D. Scott Samuels and Justin D. Radolf
c. 630 pp., March 2010

ISBN: 978-1-904455-58-5 \$310 / £159

Written by renowned scientists in the field who have made seminal contributions to the field, this book is a comprehensive guide to the pathogenic *Borrelia*.

Influenza

Molecular Virology

Edited by: Qinghua Wang and Yizhi Jane Tao
c. 200 pp., February 2010

ISBN: 978-1-904455-57-8 \$310 / £159

NS1, hemagglutinin, nucleoprotein, glycoproteins, M2 channel, virulence, polymerase, microarrays, vaccine design.

RNA Interference and Viruses

Current Innovations and Future Trends

Edited by: Miguel Angel Martínez
c. 280 pp., February 2010

ISBN: 978-1-904455-56-1 \$310 / £159

Expert RNAi specialists from around the world have teamed up to produce a timely and thought-provoking review of the area.

Retroviruses

Molecular Biology, Genomics and Pathogenesis

Edited by: Reinhard Kurth and Norbert Bannert
c. 520 pp., January 2010

ISBN: 978-1-904455-55-4 \$310 / £159

Genomics, molecular biology and pathogenesis, comprehensively covering all the recent advances.

Metagenomics

Theory, Methods and Applications

Edited by: Diana Marco
x + 212 pp., January 2010

ISBN: 978-1-904455-54-7 \$310 / £159

Essential reading for all researchers performing metagenomics studies. Highly recommended.

Aspergillus

Molecular Biology and Genomics

Edited by: M. Machida and K. Gomi
x + 238 pp., January 2010

ISBN: 978-1-904455-53-0 \$310 / £159

Systematics, bioinformatics, systems biology, regulation, genetics, genomics, metabolism, ecology, development.

Environmental Molecular Microbiology

Edited by: Wen-Tso Liu and Janet K. Jansson
viii + 232 pp., January 2010

ISBN: 978-1-904455-52-3 \$310 / £159

Current technology and applications. Microbial diversity, phylogeny, communities, 16S rRNA, metagenomics, metaproteomics, microarrays, fingerprinting, soil, water, plants, humans, biofilms.

Neisseria

Molecular Mechanisms of Pathogenesis

Edited by: Caroline Genco and Lee Wetzler
x + 270 pp., January 2010

ISBN: 978-1-904455-51-6 \$310 / £150

Genomics, biofilms, adhesion, invasion, immunity, complement, apoptosis, vaccine, epidemiology, antibiotic resistance.

Frontiers in Dengue Virus Research

Edited by: K.A. Hanley and S.C. Weaver
viii + 304 pp., January 2010

ISBN: 978-1-904455-50-9 \$310 / £150

Evolution, epidemiology, translation, replication, pathogenesis, host, animal models, mosquito interactions, transmission, vaccine, drugs, immunotherapy.

ABC Transporters in Microorganisms

Edited by: Alicia Ponte-Sucre
xii + 260 pp., August 2009

ISBN: 978-1-904455-49-3 \$310 / £150

Pili and Flagella

Current Research and Future Trends

Edited by: Ken Jarrell

x + 238 pp., August 2009

ISBN: 978-1-904455-48-6 \$310 / £150

Lab-on-a-Chip Technology

Edited by: K. E. Herold and A. Rasooly

Vol 1: Fabrication and Microfluidics

ISBN: 978-1-904455-46-2 \$310 / £150

Vol 2: Biomolecular Separation and Analysis

ISBN: 978-1-904455-47-9 \$310 / £150

Bacterial Polysaccharides

Current Research and Future Trends

Edited by: Matthias Ullrich

xii + 358 pp., June 2009

ISBN: 978-1-904455-45-5 \$310 / £150

Microbial Toxins:

Current Research and Future Trends

Edited by: Thomas Proft

viii + 192 pp., May 2009

ISBN: 978-1-904455-44-8 \$310 / £150

Acanthamoeba:

Biology and Pathogenesis

Author: Naveed Khan

viii + 290 pp., February 2009

ISBN: 978-1-904455-43-1 \$310 / £150

Bacterial Secreted Proteins: Secretory Mechanisms and Role in Pathogenesis

Edited by: Karl Wooldridge

xii + 512 pp., April 2009

ISBN: 978-1-904455-42-4 \$310 / £150