

# ***Saccharomyces boulardii* effects on gastrointestinal diseases**

**Galliano Zanello, François Meurens, Mustapha Berri, and Henri Salmon\***

Institut National de la Recherche Agronomique (INRA), UR1282, Infectiologie Animale et Santé Publique, F-37380, Nouzilly (Tours), France.

## **Abstract**

Health benefits attributed to probiotics have been described for decades. They include the treatment and the prevention of gastrointestinal diseases, vaginal and urinary infections and allergies. *Saccharomyces boulardii*, a species of yeast widely distributed, has been described as a biotherapeutic agent since several clinical trials displayed its beneficial effects in the prevention and the treatment of intestinal infections and in the maintenance of inflammatory bowel disease. All these diseases are characterized by acute diarrhoea. Administration of the yeast in combination or not with an antibiotherapy has shown to decrease significantly the duration and the frequency of diarrhoea. Experimental studies elucidated partially the molecular mechanisms triggered to improve the host health. The discovery of its anti-inflammatory and immuno-modulatory activities in correlation with the advances in the understanding of mucosal immunology opens a new field of perspectives in *S. boulardii* therapeutic applications.

## **Introduction**

The intestine contains a complex and dynamic microflora including more than 2000 micro-organism species coexisting in a complex equilibrium with the host. This microflora has various effects including metabolic activities, trophic effects on the intestinal epithelium, interactions with the host immune system (Guarner and Malagelada, 2003) and acts as a barrier to prevent colonization by opportunistic and pathogenic micro-organisms (Vollaard and Clasener, 1994). The immune system and particularly the gut-associated lymphoid tissues (GALT) provides the host with protective mechanisms against pathogen invasion across the intestinal mucosal surface (Acheson and Luccioli, 2004). However, disequilibrium in the gut microflora ecology and in the immune response could induce gastrointestinal diseases.

The term "probiotic" has been firstly defined by Fuller as "a live microbial feed supplement which beneficially affects the host by improving its intestinal microbial balance" (Fuller, 1989). This definition has been extended to health and probiotics were redefined as "live micro-organisms that, when administered in adequate amounts, confer a health benefit to the host" (FAO/WHO, 2001). *Saccharomyces Boulardii*, isolated from litchi fruit by Henri Boulard in the 1920s, belongs to the *Saccharomyces* genus which

is commonly used in several food processes including beverages and bread fermentation. This yeast is frequently prescribed in a lyophilized form as a biotherapeutic agent (Elmer et al., 1996; Klein et al., 1993). Indeed, controlled clinical trials have shown that oral administration of *S. boulardii* could treat or prevent gastrointestinal diseases such as antibiotic-associated diarrhoea (Kotowska et al., 2005), recurrent *Clostridium difficile*-associated diseases (McFarland, 2006), traveller's diarrhoea (McFarland, 2007), children acute diarrhoea (Htwe et al., 2008), enteral tube feeding-associated diarrhoea (Bleichner et al., 1997), AIDS-associated diarrhoea (Saint-Marc et al., 1991), intestinal bowel disease such as Crohn's disease and ulcerative colitis (Guslandi et al., 2003; Guslandi et al., 2000) and irritable bowel syndrome (Maupas et al., 1983).

*S. boulardii* and *S. cerevisiae* are members of the same species (Edwards-Ingram et al., 2007) but they differ genetically, metabolically and physiologically (Fietto et al., 2004; Hennequin et al., 2001). *S. boulardii* is characterized by a specific microsatellite allele (Hennequin et al., 2001) and recent studies showed that *S. boulardii* genome presents trisomy of the chromosome 9, altered copy numbers of genes potentially contributing to the increased growth rate and a better survival in acidic environment (Edwards-Ingram et al., 2007). In addition to a better resistance to acidic stress, *S. boulardii* grows faster than *S. cerevisiae* at 37°C (Fietto et al., 2004). Pharmacokinetic studies have shown that after oral administration of lyophilized *S. boulardii*, the steady-state concentrations are achieved in the colon within 3 days, and the yeast are cleared from the stools within 2-5 days after discontinuation (Blehaut et al., 1989). *S. boulardii* displays important characteristics allowing a micro-organism to transit through the gastrointestinal tract and to be used as a probiotic. During the intestinal transit, *S. boulardii* interacts with resident microflora and intestinal mucosa. Moreover, experimental studies displayed that *S. boulardii* induces a protection against enteric pathogens (Czerucka and Rampal, 2002; Mumy et al., 2007), modulates the host immune response (Ozkan et al., 2007; Rodrigues et al., 2000), decreases inflammation (Lee et al., 2005; Sougioultzis et al., 2006) and hydroelectrolytic secretions (Czerucka and Rampal, 1999), inhibits bacterial toxins (Castagliuolo et al., 1999; Tasteyre et al., 2002) and enhances trophic factors such as brush border membrane enzymes and nutrient transporters (Buts et al., 1986; Buts et al., 1994).

The aim of this review is to summarize the current knowledge about the beneficial effects of *S. boulardii* on gastrointestinal diseases. Firstly, clinical trials will be succinctly reviewed. Then, the experimental studies which have lead to a better understanding of the mechanisms used by the yeast to prevent and/or treat gastrointestinal diseases will be extensively described.

\*Corresponding author: Email: [salmon@tours.inra.fr](mailto:salmon@tours.inra.fr).  
Phone: +(33)-247427891; Fax: +(33)-247427779.

## Lactobacillus Molecular Biology: From Genomics to Probiotics | Book

Buy now!

**Publisher:** Caister Academic Press

**Editors:** Åsa Ljungh and Torkel Wadström *Lund University, Faculty of Medicine, Sweden*

**Publication date:** January 2009 [Available now!](#)

**ISBN:** 978-1-904455-41-7

**Price:** GB £150 or US \$310 (hardback).

**Pages:** x + 206

*Lactobacillus* is a genus of Gram-positive facultative anaerobic or microaerophilic bacteria. In humans they are symbiotic and are found in the gut flora. *Lactobacillus* species are used for the production of yogurt, cheese, sauerkraut, pickles, beer, wine, cider, kimchi, chocolate and other fermented foods, as well as animal feeds such as silage. In recent years much interest has been shown in the use of lactobacilli as probiotic organisms and their potential for disease prevention in humans and animals.

This major new work focuses on recent research on the molecular biology and genomics of *Lactobacillus*. Written by an international team of scientists the volume is an essential reference for all medical researchers, dairy technologists, microbiologists and biotechnologists in the academic and industrial sectors. Topics covered include phylogenetics, taxonomy, comparative genomics, functional genomics, the intestinal microflora, surface proteins, stress responses, interaction with the immune system, probiotics, anti-cancer potential, and much more. Essential reading for all scientists involved with lactic acid bacteria or probiotic research and a recommended book for all microbiology laboratories.

### Reviews:

"... an interesting blend of fundamental and applied topics relevant to the use of these important organisms in research and industry. Fundamental aspects covered in the book are taxonomy, metabolism, stress response, genomics, and surface proteins of lactobacilli. Also included are chapters on applications of *Lactobacillus* strains and their potential as probiotics in the treatment of diseases such as cancer or urinary tract infections." *from Biotechnol. J. (2009)*

"... contributions from respected international scientists, many of which are leaders in their respective fields, this book constitutes an authoritative resource about both fundamental research and applications of lactobacilli. ... essential and up to date information for anyone interested in the biology of lactobacilli. The book will clearly be of interest to microbiologists, nutritionists, food scientists, and medical practitioners alike, and it is a valuable contribution to the probiotic literature." *from Biotechnol. J. (2009)*

"*Lactobacillus Molecular Biology. From Genomics to Proteomics* is an essential reference for medical researchers, microbiologists and biotechnologists, especially, dairy technologists." *from Mercedes Berlanga, University of Barcelona writing in Int. Microbiol. (2008) 11: 294-295.*

"the chapters are edited competently and constitute outstanding independent reviews ... very informative and provides a comprehensive overview of the genus *Lactobacillus*" *from Knut Heller, Kiel writing in Biospektrum (2009) 15: 348.*

"intended for medical professionals, but the conclusions are exciting enough to convince anyone to eat more yogurt." *from SciTech Book News June 2009 p. 66*

"... a useful guide to how microbiology is developing ... a wide range of important topics ... production is beautiful, with clear diagrams, a nice typeface and two text columns per 17x25 cm page ... a most valuable text" *from Microbiology Today 2009*

### 1. Effects of *Saccharomyces boulardii* on gastrointestinal diseases: Overview in clinical trials

Several clinical trials have been performed to assess the efficacy of *S. boulardii* in the prevention and the treatment of gastrointestinal diseases which induce diarrhoea (Kotowska et al., 2005; McFarland, 2006; McFarland et al., 1994; Szajewska et al., 2007). Diarrhoea is defined as frequent watery stools resulting from impaired fluid and electrolyte absorption in the intestinal lumen, and is usually caused by pathogenic micro-organisms (Roy et al., 1995). These studies revealed the anti-diarrhoeal effects of the yeast which reduce the duration and the frequency of diarrhoea after oral administration.

#### Antibiotic-associated diarrhoea

Antibiotic-associated diarrhoea (AAD) is an acute inflammation of the intestinal mucosa caused by the administration of antibiotics and resulting in the disruption of the intestinal microflora. *Clostridium difficile* infection caused 10-20% of AAD and 95% of pseudomembranous colitis (Katz et al., 1996; Kelly et al., 1994). Other infectious organisms causing AAD include *Clostridium perfringens*, *Staphylococcus aureus*, *Klebsiella oxytoca*, *Candida* species and *Salmonella* species (Hogenauer et al., 1998). The risk factors to develop AAD are multiple: Host factors, antibiotherapy and exposure to nosocomial pathogens (Ackermann et al., 2005; McFarland, 1998). Several meta-analyses were performed to evaluate the efficacy of probiotics for the prevention of AAD (Cremonini et al., 2002; D'Souza et al., 2002; McFarland, 2006; Szajewska et al., 2006). These studies concluded that *S. boulardii* is effective in preventing AAD in adults and children. *S. boulardii* preventive effect on AAD was evaluated in 269 children with otitis media and/or respiratory tract infections (Kotowska et al., 2005). Children received antibiotic treatment plus 250 mg of *S. boulardii* (experimental group,  $n = 132$ ) or a placebo (control group,  $n = 137$ ) orally and twice a day during the antibiotherapy. Analyses included data from 246 children and demonstrated that the prevalence of diarrhoea was lower in experimental than in placebo groups (8% versus 23%, relative risk: 0.3, 95% confidence interval: 0.2-0.7). Moreover, in a recent study conducted on 151 hospitalized patients, the subjects were divided in two groups of 78 and 73 patients which received respectively antibiotics plus placebo and antibiotics plus *S. boulardii* twice a day (Can et al., 2006). The diarrhoea occurrence was investigated and the presence of *Clostridium difficile* toxin A in stools was evaluated by enzyme-linked immuno sorbent assay. The results showed a decrease in diarrhoea occurrence (1.4% versus 9% in the placebo group,  $P < 0.005$ ). In addition, *C. difficile* toxin A assay revealed 2 positive stools samples (2/7) from patients in the placebo group while nothing was detected in *S. boulardii*-treated group (0/7) (Can et al., 2006).

#### Recurrent *Clostridium difficile*-associated diseases

Treatment of *Clostridium difficile*-associated diseases with metronidazole or vancomycin is effective, but in few patients, the disease recurs after the antibiotherapy. Several clinical trials have shown the efficacy of *S. boulardii* in combination with antibiotics for treating relapses of diarrhoea and colitis (McFarland, 2006; McFarland et al., 1994; Surawicz et al., 2000). In a first randomized placebo-controlled study, 124

patients treated with antibiotics plus *S. boulardii* (500 mg twice daily) had a reduction in occurrence of diarrhoea or colitis (34.6% versus 64.7% relapses in the placebo group,  $P = 0.04$ ) (McFarland et al., 1994). Another study conducted in 170 patients showed the efficacy of *S. boulardii* (1g/d for 28 days) plus vancomycin (2g/d for 10 days) in the decrease of recurrence (16.7% versus 50% in the placebo group,  $P = 0.05$ ) (Surawicz et al., 2000). This result was significant only in patients treated with high-dose of vancomycin (2g/d) and not with lower dose (500mg/d) or metronidazole (1g/d). A recent meta-analysis (McFarland, 2006) compared the efficacy of probiotics (*S. boulardii*, *Lactobacillus rhamnosus* GG and probiotics mixtures) in the prevention of AAD and the treatment of *Clostridium difficile*-diseases. The authors concluded that the three types of probiotics significantly reduced the development of AAD but only *S. boulardii* was shown to be effective for *Clostridium difficile*-associated diseases.

#### Traveller's diarrhoea

Traveller's diarrhoea is the most frequent disorder encountered by persons travelling from low risk regions to developing areas where enteric infection is hyperendemic. Enterotoxigenic *E. coli*, *Shigella* and *Salmonella* account for about 80% of acute traveller's diarrhoea with an identified pathogen (Sanders and Tribble, 2001). In a placebo controlled double-blind study, *S. boulardii* preventive effect on traveller's diarrhoea was evaluated on a total of 1016 travellers assigned in 3 groups (Kollaritsch et al., 1993). The first group received a placebo and the second and the third groups received *S. boulardii* (250 and 1000 mg/d, respectively). Treatment with the yeast has been assumed 5 days before the travel and during the entire trip. The incidence of diarrhoea was 39.1% in the placebo group, 34.4% in the second group ( $P = 0.019$  versus placebo), and 28.7% in the third group ( $P = 0.005$ ). Recently, a meta-analysis was performed to assess the efficacy of probiotics in the prevention of traveller's diarrhoea and concluded that *S. boulardii* and a mixture of *Lactobacillus acidophilus* plus *Bifidobacterium bifidum* had a significant efficacy (McFarland, 2007).

#### Acute diarrhoea in children

Diarrhoeal diseases are a leading cause of childhood morbidity and mortality in developing countries. The aetiology of acute diarrhoea in children includes infections of the gastrointestinal tract due to bacteria, viruses or protozoa, intoxications, systemic infections, malabsorption disorders, nutritional deficiency, allergy and intolerance to food or drugs (Guandalini, 2000). There are several randomized placebo-controlled studies showing the efficacy of *S. boulardii* in the management and prevention of acute childhood diarrhoea (Htwe et al., 2008; Szajewska et al., 2007). A meta-analysis based on 5 randomized-controlled trials (619 participants) displayed this effect (Szajewska et al., 2007). Combined data showed that *S. boulardii* reduces significantly diarrhoea duration and the risk of prolonged diarrhoea compared with control. A recent study conducted in 100 hospitalized children showed that *S. boulardii* treatment for 5 days reduces the mean duration of diarrhoea (3.08 versus 4.68 days in placebo group,  $P < 0.05$ ), the frequency of stools (on day 2: 54% versus 30% in placebo group had less than 3 stools/d,  $P = 0.019$ ) and



normalizes stool consistency (on day 3: 76% *versus* 24% in placebo group,  $P = 0.019$ ) (Htwe et al., 2008).

#### **Diarrhoea in patients with total enteral feeding**

Diarrhoea is the most frequent complication in enteral tube feeding (Whelan et al., 2001). Alterations in the colonic microflora have been identified in patients receiving enteral tube feeding and these changes may be associated with the diarrhoea incidence (Schneider et al., 2000). *S. boulardii* can be used to prevent these negative alterations and to reduce the diarrhoea incidence. The preventive effect of *S. boulardii* has been assessed in a multicenter, randomized, double-blind placebo-controlled study (Bleichner et al., 1997). A total of 128 critically ill patients needing enteral nutrition for at least 6 days were included in the study. *S. boulardii* administered at the dose of 2 g/d reduced the mean percentage of days with diarrhoea (14.2% *versus* 18.9% in the placebo group, odds ratio = 0.67, 95%, confidence interval: 0.50-0.90,  $P = 0.0069$ ). This study confirmed the results obtained in two previous clinical trials where *S. boulardii* administration decreased the length of diarrhoea events (Schlotterer et al., 1987; Tempe et al., 1983).

#### **AIDS-associated diarrhoea**

Acquired immune deficiency syndrome (AIDS) is a viral infection characterized by immune cell dysfunction and subsequent immunodeficiency, as well as intestinal disorder (Kotler et al., 1984). A randomized, double-blind placebo-controlled study conducted in 35 patients showed that administration of *S. boulardii* (3 g/d) for one week decreases the diarrhoea incidence. After 7 days of treatment, 61% of patients were diarrhoea-free compared with 12% in the placebo group ( $P < 0.002$ ) (Saint-Marc et al., 1991).

#### **Inflammatory bowel diseases**

Inflammatory bowel diseases (IBD) are characterized by a chronic course in which phases of remission of variable length are interrupted by acute episodes. The two main subforms are ulcerative colitis and Crohn's disease. Intestinal inflammation in ulcerative colitis is primarily limited to the colon whereas the whole gastrointestinal tract can be involved in Crohn's disease (Xavier and Podolsky, 2007). In IBD, genetically defects result in abnormal mucosal immune response overacting the microflora and inducing intestinal inflammation (Strober et al., 2007). In a pilot study conducted in 20 patients, administration of *S. boulardii* in addition of conventional therapy was found superior to placebo in promoting reduction of bowel movements (Plein and Hotz, 1993). Furthermore, in another clinical trial conducted in 32 patients, the preventive effect of *S. boulardii* was evaluated in relapses of Crohn's disease (Guslandi et al., 2000). After six months, patients treated with mesalamine alone (3 g/d) presented more clinical relapses compared to patients receiving *S. boulardii* (1 g/d) plus mesalamine (2 g/d) (37.5% *versus* 6.25%,  $P = 0.04$ ). The same author evaluated the effectiveness of *S. boulardii* in the treatment of ulcerative colitis in 25 patients. Administration of *S. boulardii* (750 mg/d) plus mesalamine for 4 weeks resulted in clinical remission for 68% of patients (Guslandi et al., 2003).

#### **Irritable bowel syndrome**

Irritable bowel syndrome (IBS) is a chronic disorder involving a combination of abdominal pain and change in

bowel habit and bloating. Recent evidence suggests a role of the microflora in IBS pathogenesis (Parkes et al., 2008). In a double-blind, placebo-controlled study conducted in 34 patients with predominant episodes of diarrhoea, treatment with *S. boulardii* decreased the daily number of stools ( $P < 0.05$ ) and improved their consistency ( $P < 0.05$ ) (Maupas et al., 1983).

#### **Safety of administration**

*S. boulardii* is administered to patients in a lyophilized form and the treatment is well tolerated. However, some rare cases of *S. boulardii* fungemias have been reported in patients with an indwelling central venous catheter (de Llanos et al., 2006; Hennequin et al., 2000; Lherm et al., 2002). The origin of the fungemia is thought to be either a digestive tract translocation or a contamination of the central venous line by the colonized hands of health workers (Hennequin et al., 2000). This raises the question of the risk-benefit ratio of *S. boulardii* in critically ill or immunocompromised patients. Thus, administration of *S. boulardii* should be contraindicated for patients of fragile health, as well as for patients with central venous catheter (Herbrecht and Nivoix, 2005).

### **2. Experimental effects of *Saccharomyces boulardii***

These effects have been assessed in several studies and showed that *S. boulardii* exerts beneficial mechanisms in animal models displaying IBD as well as in pathogenic or opportunistic micro-organism infection models such as *Clostridium difficile*, *Vibrio cholerae*, *Escherichia coli*, *Salmonella enterica* subspecies *enterica* serovar Typhimurium, *Shigella flexneri*, *Citrobacter rodentium* and *Candida albicans* (Chen et al., 2006; Czerucka et al., 1994; Dahan et al., 2003; Dalmasso et al., 2006b; Jawhara and Poulain, 2007; Mumy et al., 2007; Wu et al., 2008). These mechanisms include the modification of host cell signalling pathways implicated in proinflammatory response and in hydroelectrolytic secretion, the stimulation of host immune defences, the neutralization of bacterial toxins and the decrease of bacterial adherence to intestinal epithelial cells, the maintenance of membrane permeability and the inhibition of pathogen translocation. Experimental effects of *S. boulardii* on diarrhoeal pathogens have been previously reviewed (Czerucka and Rampal, 2002).

#### **2.1. Anti-inflammatory effects**

Several experimental studies showed that *S. boulardii* interferes with the host cell signalling pathways and decreases the expression of inflammation-associated cytokines such as interleukin 8 (IL-8), IL-6, IL-1 $\beta$ , tumor necrosis factor alpha (TNF- $\alpha$ ) and interferon gamma (IFN- $\gamma$ ) (Dahan et al., 2003; Dalmasso et al., 2006b; Mumy et al., 2007). These studies showed that the yeast can reduce inflammation in blocking nuclear factor-kappa B (NF- $\kappa$ B) (Dahan et al., 2003; Mumy et al., 2007) and mitogen-activated protein kinase (MAPK) activation (Chen et al., 2006; Mumy et al., 2007), in decreasing nitric oxide (NO) production (Girard et al., 2005), in enhancing peroxisome proliferators-activated receptor-gamma (PPAR- $\gamma$ ) expression (Lee et al., 2005) and in modulating T cell migratory behaviour (Dalmasso et al., 2006a).

#### *Inhibition of NF- $\kappa$ B and MAPK activation*

During the course of infection, activation of NF- $\kappa$ B and MAPK in intestinal epithelial cells results in pro-inflammatory cytokines secretion and intestinal inflammation. *Clostridium difficile*-toxin A induces intestinal inflammation, fluid secretion and mucosal injury (Torres et al., 1990; Triadafilopoulos et al., 1987; Triadafilopoulos et al., 1989). After stimulation with toxin A, *S. boulardii* supernatant (<10 kDa) has shown to decrease IL-8 secretion in human colonocytes and extracellular signal-regulated protein kinases 1 and 2 (ERK1/2) activation in both human colonocytes and murine ileal loops (Chen et al., 2006). In a model of T84 human colonic cell line, *S. boulardii* inhibited also the NF- $\kappa$ B and MAPK signalling pathways in enterohemorrhagic *E. coli* (EHEC) infected-T84 cells resulting in the decrease of IL-8 and TNF- $\alpha$  secretion by epithelial cells (Dahan et al., 2003; Dalmasso et al., 2006b). Furthermore, the decrease of IL-8 secretion was observed in *Shigella flexneri* infected-T84 cells when the yeast or the yeast conditioned medium were added to the monolayer culture (Mumy et al., 2007). This decrease was greater for cells treated with the conditioned medium. Furthermore, polymorphonuclear leukocytes (PMN) transepithelial migration is reduced in presence of the yeast. This result could be explained by the decrease in IL-8 secretion level since this cytokine has been previously shown to be chemoattractant for PMN. *S. boulardii* decreased also phosphorylation and activation of ERK and JNK MAP Kinases and decreased the I $\kappa$ B phosphorylation and degradation during the infection, thereby preventing NF- $\kappa$ B activation (Mumy et al., 2007). These events are likely responsible for the decrease in IL-8 secretion and PMN transmigration. These anti-inflammatory effects were confirmed *in vivo* with a model of human fetal colonic tissue transplanted into severe combined immunodeficiency (SCID) mice (Mumy et al., 2007). In addition, recent data showed that the yeast produces a low molecular weight soluble factor (< 1 kDa) blocking NF- $\kappa$ B activation and NF- $\kappa$ B-mediated IL-8 gene expression in intestinal epithelial cells and monocytes (Sougioultzis et al., 2006) (Figure 1).

#### *Decrease of nitric oxide production*

In IBD, production of high levels of NO is associated with inflammatory effects (Dijkstra et al., 1998). NO is released through conversion of L-arginine to NO and L-citrulline. This reaction is catalyzed by three isoforms of the nitric oxide synthase (NOS): Neuronal NOS, endothelial NOS and inducible NOS (Moncada et al., 1991). The iNOS activity is up-regulated during immune activation and can result in the synthesis of high levels of NO. *S. boulardii* inhibitory effect in iNOS activity was displayed in the rat castor oil-induced diarrhoea model (Girard et al., 2005). The citrulline level (a marker of NO production) was increased in the colon of rats whereas *S. boulardii* administration was shown to block this production (Figure 1). The same effect is reported with the iNOS inhibitor. These results suggest that iNOS inhibition by *S. boulardii* may be beneficial in the treatment of diarrhoea and/or IBD associated with overproduction of NO (Girard et al., 2005).

#### *Enhancement of PPAR- $\gamma$ expression*

PPAR- $\gamma$  is a nuclear receptor expressed by several cell types including intestinal epithelial cells, dendritic cells (DC), T and B cells, and can act as a regulator of the inflammation

(Escher et al., 2001; Mansen et al., 1996; Su et al., 1999). It has been shown that *S. boulardii* up-regulates PPAR- $\gamma$  expression in human colonocytes at both mRNA and protein levels (Lee et al., 2005). This up-regulation is correlated with a reduction of the human colon cell response to proinflammatory cytokines (Lee et al., 2005) (Figure 1). This anti-inflammatory effect was supported by previous data showing that PPAR- $\gamma$  ligands inhibited NF- $\kappa$ B activation and that TLR4-activated PPAR- $\gamma$  in epithelium attenuated NF- $\kappa$ B signalling (Dubuquoy et al., 2003; Genolet et al., 2004). A recent study showed that PPAR- $\gamma$  expression in colonic epithelial cells decreased pro-inflammatory cytokines levels and protects mice against experimental IBD (Adachi et al., 2006). Thus, *S. boulardii* up-regulates PPAR- $\gamma$  expression which could be a therapeutic target in IBD.

#### *Modulation of T cell migratory behaviour*

In a model of lymphocyte-transferred SCID mice displaying IBD, *S. boulardii* administration decreased the production of IFN- $\gamma$  and inflammation in the colon (Dalmasso et al., 2006a). This observation was correlated with a modification of T cells distribution. Indeed, the number of IFN $\gamma$ -producing CD4+ T cells decreased within the colon and there were more cells in the mesenteric lymph node. *S. boulardii* appeared to be involved in the modification of lymph node endothelial cell adhesiveness and in the recruitment of IFN $\gamma$ -producing CD4+ T cells to this location. These results suggested that *S. boulardii* administration could have indirectly beneficial effect in the treatment of IBD (Dalmasso et al., 2006a).

## **2.2. Immuno-modulatory effects**


Among the mechanisms used by probiotics to improve the host health, one of them is the stimulation of the host immunity. Several studies demonstrated that *S. boulardii* might induce a protective effect in modulating both innate and adaptive host immunity to respond against pathogen infection.

#### *Saccharomyces boulardii effect on innate immunity*

*S. boulardii* immunopharmacological effects in healthy human volunteers have been evaluated and it has been shown that the yeast triggered the activation of the complement and the reticuloendothelial systems (Caetano et al., 1986). *In vitro*, the yeast activates the complement and the migration of monocyte and granulocyte (Caetano et al., 1986). Furthermore, *S. boulardii* administration to germ-free mice has shown to enhance the number of K $\ddot{u}$ pffer cells (Rodrigues et al., 2000). These results indicate that *S. boulardii* is able to stimulate the innate immune system.

#### *Saccharomyces boulardii effect on adaptive immunity*

*S. boulardii* oral administration to rats and mice has shown to stimulate the secretion of immune factors (Buts et al., 1990; Qamar et al., 2001; Rodrigues et al., 2000). In the duodenal fluid of rats treated with the yeast, secretory immunoglobulin A (sIgA) mean concentration was increased by 56.9% (*versus* control,  $P < 0.01$ ) (Buts et al., 1990). Additionally, the secretory component was significantly increased in the duodenal fluid (62.8% *versus* control,  $P < 0.01$ ) as well as in villus cells (69% *versus* control,  $P < 0.025$ ) and in crypt cells (80% *versus* control,  $P < 0.01$ ) (Buts et al., 1990). The same effect was reported in mice challenged with *C. difficile* toxin A (Qamar et al., 2001). Treatment with *S. boulardii*



**Figure 1.** During the course of infection with enteric pathogens such as *Clostridium difficile*, *Escherichia coli* or *Shigella flexneri*, activation of nuclear factor-kappa B (NF-κB) and mitogen-activated protein kinase (MAPK) in enterocyte leads to the secretion of inflammation associated cytokines such as IL-8 and TNF-α. *Saccharomyces boulardii* (Sb) secretes soluble factors which reduce NF-κB and MAPK activation resulting in a decrease of the inflammation. The yeast produces a low molecular weight soluble factor (< 1 kDa) blocking NF-κB-mediated IL-8 expression. In inflammatory bowel disease (IBD), production of high levels of nitric oxide (NO) is correlated with intestinal inflammation. *Saccharomyces boulardii* decreases the inducible nitric oxide synthase (iNOS) activity and consequently reduces intestinal inflammation. In IBD, inhibition of peroxisome proliferators-activated receptor-gamma (PPAR-γ), a regulator of inflammation, leads to intestinal inflammation. *Saccharomyces boulardii* up-regulates PPAR-γ expression and consequently decreases intestinal inflammation.

caused a 1.8-fold increase in total sIgA levels ( $P = 0.003$ ) and a 4.4-fold increase in specific anti-toxin A sIgA levels ( $P < 0.001$ ). The effect of *S. boulardii* on the immune system was also evaluated by comparison of germ-free mice mono-associated with the probiotic and germ-free mice non-


associated (Rodrigues et al., 2000). Higher levels of sIgA both total and anti-*S. boulardii* were produced in the small intestine of mono-associated mice (Rodrigues et al., 2000). Thus, these results demonstrated that the yeast is able to enhance the secretion of mucosal immune factors which protect the host against enteric pathogens (Figure 2A).

After intravenous administration of *E. coli* B41, clearance of the human pathogenic strain from the bloodstream of *S. boulardii* monoassociated mice was more effective than in germ-free mice. More efficient clearance was correlated with earlier production of IFN- $\gamma$  and IL-12 which are involved in the T-helper 1 response (Rodrigues et al., 2000). A further study in healthy human volunteers investigated the effect of *S. boulardii* on lymphocyte phenotype (Jahn et al., 1996). Results were compared before and after the yeast treatment. No phenotypic changes were observed in intestinal lymphocytes. However, CD4<sup>+</sup> T cells of the peripheral blood had a significant increased expression of CD25 ( $P < 0.02$ ) (Jahn et al., 1996). Since it has been shown that CD4<sup>+</sup> CD25<sup>+</sup> T cells can display regulatory functions (Sakaguchi et al., 1995; Sakaguchi et al., 2007), further studies should be performed to assess the effect of *S. boulardii* in the stimulation of regulatory T cell population and to investigate a possible link between the induction of a regulatory immune response and the decrease of inflammation.

### 2.3. Inhibition of hydroelectrolytic secretions

*S. boulardii* inhibition of hydroelectrolytic secretions was widely demonstrated in *Vibrio cholerae* infectious model (Czerucka et al., 1989; Czerucka et al., 1994; Vidon et al., 1986). *Vibrio cholerae* produces a 84 kDa toxin (CT) composed of two subunits: A and B. CT binds its intestinal receptor (monosialoganglioside GM1) through the B subunit and catalyzes the activation of adenylate cyclase through the A subunit. This activation results in an increase on cyclic adenosine monophosphate (cAMP) levels triggering active secretion of chloride and bicarbonate in crypt cells and inhibiting chloride absorption in the villi. This perturbation in hydroelectrolytic secretions induces severe diarrhoea (Kaper et al., 1994).

In a previous study, the model of jejunal intestinal loops in rats allowed to demonstrate the protective effect of *S. boulardii* in CT-induced pathogenesis (Vidon et al., 1986). The loops injected with CT and then with the yeast presented a significant decrease in the hydrosaline secretion. The



**Figure 2.** A: Oral administration of *Saccharomyces boulardii* (Sb) enhances both secretory IgA (sIgA) concentration in the intestinal lumen and brush border membrane enzyme (BBM) expression. The increase in their expression seems to be correlated with a secretion of polyamines during the yeast catabolism.

B: Cholera toxin (CT) binding on the intestinal epithelial cell surface induces adenylate cyclase (AC) expression resulting in chloride (Cl<sup>-</sup>) secretion. *Saccharomyces boulardii* produces a 120 kDa protein which inhibits AC activation resulting in a decrease of chloride secretion.

C: *Saccharomyces boulardii* neutralizes bacterial toxins. The yeast produces respectively a 54 kDa proteinase which exerts a proteolytic activity on both *Clostridium difficile* toxins A and B and a 63 kDa protein phosphatase which dephosphorylates LPS from enteropathogenic *Escherichia coli*.

D: Enteric pathogens such as enterohemorrhagic (EHEC) and enteropathogenic (EPEC) *Escherichia coli* and *Shigella flexneri* induce an alteration of the epithelial barrier integrity in affecting the tight-junctions-associated proteins. *Saccharomyces boulardii* abolishes the myosin light chain (MLC) phosphorylation and restores *Zonula occludens* 1 and 2 (ZO-1, ZO-2) and claudin-1 distribution. Therefore, the yeast enhances the cell ability to restore intestinal permeability.



inhibitory effect of *S. boulardii* on CT was confirmed using rats' intestinal cell lines (IRD-98 and IEC-17) (Czerucka et al., 1989). Cells pre-incubated with the yeast and challenged with the CT had a 50% decrease in the activity of cAMP. However, in this study, when *S. boulardii* was killed, the inhibitory effect on cAMP disappeared. The same author showed that incubation of IEC-6 cells (rat intestinal epithelial cells) with the yeast-conditioned medium decrease the cAMP activity induced by CT. This effect disappeared when the yeast-conditioned medium was denatured by heating, by trichloroacetic acid precipitation or by trypsin hydrolysis suggesting the presence of a secreted factor which has been characterized as a 120 kDa protein (Czerucka et al., 1994) (Figure 2B). These results drove Czerucka and Rampal to evaluate in T84 cells the effect of *S. boulardii*-conditioned medium (Sb-CM) on chloride secretion mediated by cAMP and calcium. Preincubation of cells with Sb-CM reduced chloride secretion mediated by cAMP agonists (CT, prostaglandin E<sub>2</sub>, vasoactive intestinal polypeptide and forskolin) and by charbachol, a muscarinic agonist implicated in the calcium signalling pathways (Czerucka and Rampal, 1999). These studies demonstrate that *S. boulardii* secretes a 120 kDa protein that interferes with chloride secretion pathways and decreases diarrhoea caused by *Vibrio cholerae* infection. An additional mechanism of protection was proposed with the adhesion of CT to a receptor on the yeast surface (Brandao et al., 1998). The authors suggested that the yeast receptor could be able to bind the B subunit while the A subunit was internalized and stimulated cAMP signals as well as cAMP-dependent trehalase activity. This result supposes that the yeast and the mammalian CT receptors could be structurally and functionally similar.

#### 2.4. Neutralization of bacterial toxins

It has been shown that *S. boulardii* has the capacity to release factors which neutralize bacterial toxins and decrease the deleterious effects of infectious pathogens (Buts et al., 2006; Castagliuolo et al., 1996; Castagliuolo et al., 1999). *C. difficile* is responsible of AAD in humans and animals, and is one of the most common nosocomial pathogens. Pathogenic *C. difficile* produces two exotoxins, toxin A and toxin B which induce intestinal inflammation, fluid secretion and mucosal injury (Pothoulakis, 1996). *S. boulardii* produces a 54 kDa serine protease exerting a proteolytic activity on both toxin A and B (Castagliuolo et al., 1996; Castagliuolo et al., 1999). This proteolysis results in the inhibition of toxins binding on their intestinal receptors (Figure 2C). Thus *S. boulardii* treatment enhances transepithelial resistance, epithelial barrier integrity and decreases hydroelectrolytic secretions in both rat ileum and colonic mucosa. *S. boulardii* produces also a 63 kDa protein phosphatase that inhibits the LPS toxicity of enteropathogenic *E. coli* O55B5 by endotoxin dephosphorylation (Buts et al., 2006) (Figure 2C). Intraperitoneal injection of dephosphorylated LPS in rats decreases the level of circulating TNF- $\alpha$  and abolishes inflammatory lesions in comparison to rats injected with native LPS.

#### 2.5. Decrease of bacterial adhesion to intestinal epithelial cells

*C. rodentium* is a micro-organism that colonizes the colon of mice, causing "attaching/effacing" lesions and colonic

hyperplasia (Schauer and Falkow, 1993). In addition to serving as model for EPEC and EHEC infection in small-animal, *C. rodentium* can also serve as model for inflammatory bowel disease in mouse (Higgins et al., 1999). Recently, the beneficial effect of *S. boulardii* on *C. rodentium*-induced colitis was assessed in mice (Wu et al., 2008). The improvement effect of *S. boulardii* was associated with significantly reduced numbers of mucosal adherent bacteria compared with infected untreated animals ( $P < 0.05$ ). This effect was not due to a bactericidal action but was correlated with a reduction in EspB and Tir protein secretions, respectively a translocator and an effector protein implicated in the type III secretion system (TTSS). The authors concluded that *S. boulardii* maintains the epithelial barrier integrity and ameliorates inflammatory lesions associated with *C. rodentium* infection by attenuating bacterial adherence to the host cells through putative actions on TTSS (Wu et al., 2008). However, in some *in vitro* models such as EPEC-infected T84 cells or *Shigella*-infected T84 cells (Czerucka 2000, Mumy 2007), *S. boulardii* did not alter the number of adherent bacteria.

#### 2.6. Maintenance of epithelial barrier integrity

Several pathogenic microorganisms such as *Shigella flexneri*, enterohemorrhagic and enteropathogenic *E. coli* have mechanisms of infection characterized by bacterial adhesion to the intestinal mucosa resulting in alteration of tight-junctions, disruption of membrane permeability and enterocyte secretion of pro-inflammatory cytokines. *S. boulardii* acts on the epithelial barrier in improving tight-junctions structure and in restoring membrane permeability disrupted by infectious pathogens (Czerucka et al., 2000; Dahan et al., 2003; Mumy et al., 2007). In EPEC E2348/69-infected T84 cells the monolayer transepithelial resistance was decreased and the perijunctional distribution of tight-junction-associated protein *Zonula occludens* (ZO-1) was altered, resulting in the disruption of epithelial barrier structure (Czerucka et al., 2000). In contrast, when the infection was performed in the presence of *S. boulardii*, the transepithelial resistance was not altered and ZO-1 protein distribution was preserved, suggesting a protective effect of *S. boulardii* on the tight-junctions structure of T84-infected cells (Czerucka et al., 2000) (Figure 2D). Moreover, the yeast prevented the caspase-3 activation induced by EPEC infection and delayed apoptosis of T84 cells. In this study, *S. boulardii* decreases EPEC induced ERK1/2-activation which seems to be correlated with the decrease of the number of intracellular bacteria (Czerucka et al., 2000). This study demonstrates that *S. boulardii* modulates the signalling pathways induced by bacterial infection and thus exerts protective effects. The same effects were observed after pre-incubation of *S. boulardii* in EHEC-infected T84 cells (Dahan et al., 2003). During the bacterial infection, the myosin light chain protein (MLC) is phosphorylated and the tight-junctions are disrupted. The MLC is associated with a cytoskeletal protein in intercellular tight-junctions control. The yeast abolished MLC phosphorylation and allowed to preserve the barrier function (Figure 2D). In *Shigella*-infected T84 cells, the yeast positively affects tight-junctions proteins (claudin-1 and ZO-2) and significantly protects the barrier function (Mumy et al., 2007) (Figure 2D). *Shigella*-infected cellular monolayer had a dramatic decrease in claudin-1 and ZO-2 levels. In the presence of the yeast, cellular



monolayer exhibited larger amounts of these proteins. After 2 hours of infection, monolayers exposed to both bacteria and yeast displayed 75% of the barrier function compared to uninfected controls (*versus* 50% for monolayers exposed to bacteria alone). These results demonstrate that *S. boulardii* enhances the cell ability to restore tight-junctions structure and barrier permeability.

### 2.7. Trophic effects on intestinal mucosa

Acute and chronic gastrointestinal diseases often induce diarrhoea. The microflora and the intestinal mucosa are disrupted, resulting in a deficit of intestinal enzymatic activity and transporter expression, and an increase of inflammation and water loss. Several studies have shown that *S. boulardii* exerts trophic effects restoring the intestinal homeostasis (Buts et al., 1986; Buts et al., 1999; Buts et al., 2002; Jahn et al., 1996; Schneider et al., 2005).

A study conducted in human volunteers and in growing rats showed that oral administration of *S. boulardii* for 8 days increased the activity of sucrase-isomaltase (+82%), lactase (+77%) and maltase-glucoamylase (+75%). This increase was also observed in growing rats. However, this treatment did not modify the intestinal morphology (Buts et al., 1986). These results were confirmed in another study reporting an increase in lactase,  $\alpha$ -glucosidase and alkaline phosphatase activities after the yeast treatment (Jahn et al., 1996). These results showed that *S. boulardii* enhances the release of brush-border membrane enzymes implicated in the nutrient degradation and absorption. These intestinal enzymes are often altered during acute or chronic enteropathies. In a further study, oral administration of the yeast during 8 days after a 60% proximal enterectomy in rats improved disaccharidase activities, increased the absorption of D-glucose coupled to  $\text{Na}^+$  as well as the expression of the sodium/glucose cotransporter-1 (SGLT-1) (Buts et al., 1999). SGLT-1 is implicated in water and electrolytes reabsorption. Thus, these trophic effects enhanced by *S. boulardii* could be efficient in the treatment of diarrhoea and congenital sucrase-isomaltase deficiency. Buts and colleagues suggested that the increased expression of intestinal enzymes could be in part due to an endoluminal release of polyamines by *S. boulardii*, mainly spermine and spermidin (Buts et al., 1994) (Figure 2A). When rats were treated with 100 mg of lyophilized *S. boulardii* containing 679 nmol of polyamines or with 500 nmol of purified spermine, similar enzymatic responses were observed with an increase in sucrase and maltase activities. Another effect of the yeast is based on the modification of luminal short-chain fatty acids (SCFAs) concentration (Schneider et al., 2005). SCFAs which are among the most important metabolites produced by anaerobic bacteria in the colon are involved in water and electrolyte absorption by the colonic mucosa (Bowling et al., 1993). Patients on long-term total enteral nutrition have a decrease in the number of fecal anaerobic bacteria and in fecal SCFAs concentration (Schneider et al., 2000). A study showed that TEN-patients treated with *S. boulardii* had an increase in total fecal SCFAs levels and total SCFAs remained high 9 days after discontinuation of the treatment (Schneider et al., 2005). The author concluded that this increase may explain the preventive effects of the yeast in TEN-induced diarrhoea. Finally, a further study demonstrated that the yeast releases a leucine aminopeptidase enhancing

endoluminal N-terminal peptide hydrolysis in suckling rat small intestine and potentially preventing reactions to food antigens when the barrier permeability is altered (Buts et al., 2002).

### Conclusion

Gastrointestinal diseases are characterized by an alteration of the microbial balance and the intestinal homeostasis. Several clinical trials and experimental studies displayed the role of *S. boulardii* as a good biotherapeutic agent allowing to prevent and/or treat several gastrointestinal diseases. In comparison to probiotic bacteria, the use of probiotic yeast is beneficial when the treatment is combined to antibiotherapy. Oral treatment with *S. boulardii* induces modulation of the host cell signalling pathways implicated in proinflammatory response and in hydroelectrolytic secretion, neutralization of bacterial toxins, inhibition of pathogen translocation, stimulation of the host immune response, restoration of intestinal permeability, and stimulation of brush-border membrane enzymes and transporters. Thus, *S. boulardii* is involved in the restoration of intestinal homeostasis. These beneficial effects are in part mediated by secreted factors such as proteases, phosphatases and polyamines.

The recent advance in mucosal immunology understanding and the discovery of *S. boulardii* immuno-modulatory and anti-inflammatory effects open a new field of perspectives. Several reports showed that *S. boulardii* induces a local mucosal protection in the gut (Buts et al., 1990; Qamar et al., 2001; Rodrigues et al., 2000). However, further studies should be performed to evaluate if the yeast can proceed as an "immunobiotic" by modulating adaptive immune factors to act in other mucosal sites than the gut as the bronchial or the uro-genital mucosa (Clancy, 2003). Another question is to identify if the yeast can activate adaptive immune factors involved in the entero-mammary link, conferring better protection to the newborn (Bourges et al., 2008). Indeed, it has been shown that the yeast can modulate the migratory behaviour of lymphocytes by modification of lymph node endothelial cell adhesiveness (Dalmasso et al., 2006a). Then, a better characterization of the pathways involved in the anti-inflammatory response and the molecular mechanisms implied is required. The discovery of T regulatory cells with anti-inflammatory functions could lead to investigate if *S. boulardii* is able to stimulate this population. In addition, *S. boulardii* effect on DCs could be investigated to identify if the yeast could stimulate DC regulatory functions by targeting specific pattern-recognition receptors.

In conclusion, *S. boulardii* activities in the prevention and/or the treatment of diarrhoea have been widely investigated and demonstrated. However, new data and further experimental studies should permit to better elucidate the mechanisms of action of the yeast and suggest new therapeutic applications.

### References

- Acheson, D. W., and Luccioli, S. (2004). Microbial-gut interactions in health and disease. *Mucosal immune responses*. *Best Pract Res Clin Gastroenterol* 18, 387-404.
- Ackermann, G., Thomalla, S., Ackermann, F., Schaumann, R., Rodloff, A. C., and Ruf, B. R. (2005). Prevalence and

- characteristics of bacteria and host factors in an outbreak situation of antibiotic-associated diarrhoea. *J Med Microbiol* 54, 149-153.
- Adachi, M., Kurotani, R., Morimura, K., Shah, Y., Sanford, M., Madison, B. B., Gumucio, D. L., Marin, H. E., Peters, J. M., Young, H. A., and Gonzalez, F. J. (2006). Peroxisome proliferator activated receptor gamma in colonic epithelial cells protects against experimental inflammatory bowel disease. *Gut* 55, 1104-1113.
- Blehaut, H., Massot, J., Elmer, G. W., and Levy, R. H. (1989). Disposition kinetics of *Saccharomyces boulardii* in man and rat. *Biopharm Drug Dispos* 10, 353-364.
- Bleichner, G., Blehaut, H., Mentec, H., and Moysse, D. (1997). *Saccharomyces boulardii* prevents diarrhea in critically ill tube-fed patients. A multicenter, randomized, double-blind placebo-controlled trial. *Intensive Care Med* 23, 517-523.
- Bourges, D., Meurens, F., Berri, M., Chevalere, C., Zanello, G., Levast, B., Melo, S., Gerdt, V., and Salmon, H. (2008). New insights into the dual recruitment of IgA(+) B cells in the developing mammary gland. *Mol Immunol* 45, 3354-3362.
- Bowling, T. E., Raimundo, A. H., Grimbale, G. K., and Silk, D. B. (1993). Reversal by short-chain fatty acids of colonic fluid secretion induced by enteral feeding. *Lancet* 342, 1266-1268.
- Brandao, R. L., Castro, I. M., Bambirra, E. A., Amaral, S. C., Fietto, L. G., Tropia, M. J., Neves, M. J., Dos Santos, R. G., Gomes, N. C., and Nicoli, J. R. (1998). Intracellular signal triggered by cholera toxin in *Saccharomyces boulardii* and *Saccharomyces cerevisiae*. *Appl Environ Microbiol* 64, 564-568.
- Buts, J. P., Bernasconi, P., Vaerman, J. P., and Dive, C. (1990). Stimulation of secretory IgA and secretory component of immunoglobulins in small intestine of rats treated with *Saccharomyces boulardii*. *Dig Dis Sci* 35, 251-256.
- Buts, J. P., Bernasconi, P., Van Craynest, M. P., Maldague, P., and De Meyer, R. (1986). Response of human and rat small intestinal mucosa to oral administration of *Saccharomyces boulardii*. *Pediatr Res* 20, 192-196.
- Buts, J. P., De Keyser, N., and De Raedemaeker, L. (1994). *Saccharomyces boulardii* enhances rat intestinal enzyme expression by endoluminal release of polyamines. *Pediatr Res* 36, 522-527.
- Buts, J. P., De Keyser, N., Marandi, S., Hermans, D., Sokal, E. M., Chae, Y. H., Lambotte, L., Chanteux, H., and Tulkens, P. M. (1999). *Saccharomyces boulardii* upgrades cellular adaptation after proximal enterectomy in rats. *Gut* 45, 89-96.
- Buts, J. P., De Keyser, N., Stilmant, C., Sokal, E., and Marandi, S. (2002). *Saccharomyces boulardii* enhances N-terminal peptide hydrolysis in suckling rat small intestine by endoluminal release of a zinc-binding metalloprotease. *Pediatr Res* 51, 528-534.
- Buts, J. P., Dekeyser, N., Stilmant, C., Delem, E., Smets, F., and Sokal, E. (2006). *Saccharomyces boulardii* produces in rat small intestine a novel protein phosphatase that inhibits *Escherichia coli* endotoxin by dephosphorylation. *Pediatr Res* 60, 24-29.
- Caetano, J. A., Parnes, M. T., Babo, M. J., Santos, A., Ferreira, A. B., Freitas, A. A., Coelho, M. R., and Mateus, A. M. (1986). Immunopharmacological effects of *Saccharomyces boulardii* in healthy human volunteers. *Int J Immunopharmacol* 8, 245-259.
- Can, M., Besirbellioglu, B. A., Avci, I. Y., Beker, C. M., and Pahsa, A. (2006). Prophylactic *Saccharomyces boulardii* in the prevention of antibiotic-associated diarrhea: a prospective study. *Med Sci Monit* 12, P119-22.
- Castagliuolo, I., LaMont, J. T., Nikulasson, S. T., and Pothoulakis, C. (1996). *Saccharomyces boulardii* protease inhibits *Clostridium difficile* toxin A effects in the rat ileum. *Infect Immun* 64, 5225-5232.
- Castagliuolo, I., Riegler, M. F., Valenick, L., LaMont, J. T., and Pothoulakis, C. (1999). *Saccharomyces boulardii* protease inhibits the effects of *Clostridium difficile* toxins A and B in human colonic mucosa. *Infect Immun* 67, 302-307.
- Chen, X., Kokkotou, E. G., Mustafa, N., Bhaskar, K. R., Sougioultzis, S., O'Brien, M., Pothoulakis, C., and Kelly, C. P. (2006). *Saccharomyces boulardii* inhibits ERK1/2 mitogen-activated protein kinase activation both *in vitro* and *in vivo* and protects against *Clostridium difficile* toxin A-induced enteritis. *J Biol Chem* 281, 24449-24454.
- Clancy, R. (2003). Immunobiotics and the probiotic evolution. *FEMS Immunol Med Microbiol* 38, 9-12.
- Cremonini, F., Di Caro, S., Nista, E. C., Bartolozzi, F., Capelli, G., Gasbarrini, G., and Gasbarrini, A. (2002). Meta-analysis: the effect of probiotic administration on antibiotic-associated diarrhoea. *Aliment Pharmacol Ther* 16, 1461-1467.
- Czerucka, D., Dahan, S., Mograbi, B., Rossi, B., and Rampal, P. (2000). *Saccharomyces boulardii* preserves the barrier function and modulates the signal transduction pathway induced in enteropathogenic *Escherichia coli*-infected T84 cells. *Infect Immun* 68, 5998-6004.
- Czerucka, D., Nano, J. L., Bernasconi, P., and Rampal, P. (1989). [Response to cholera toxin of 2 epithelial intestinal cell lines. Effect of *Saccharomyces boulardii*]. *Gastroenterol Clin Biol* 13, 383-387.
- Czerucka, D., and Rampal, P. (1999). Effect of *Saccharomyces boulardii* on cAMP- and Ca<sup>2+</sup>-dependent Cl<sup>-</sup> secretion in T84 cells. *Dig Dis Sci* 44, 2359-2368.
- Czerucka, D., and Rampal, P. (2002). Experimental effects of *Saccharomyces boulardii* on diarrheal pathogens. *Microbes Infect* 4, 733-739.
- Czerucka, D., Roux, I., and Rampal, P. (1994). *Saccharomyces boulardii* inhibits secretagogue-mediated adenosine 3',5'-cyclic monophosphate induction in intestinal cells. *Gastroenterology* 106, 65-72.
- D'Souza, A. L., Rajkumar, C., Cooke, J., and Bulpitt, C. J. (2002). Probiotics in prevention of antibiotic associated diarrhoea: meta-analysis. *BMJ* 324, 1361.
- Dahan, S., Dalmasso, G., Imbert, V., Peyron, J. F., Rampal, P., and Czerucka, D. (2003). *Saccharomyces boulardii* interferes with enterohemorrhagic *Escherichia coli*-induced signaling pathways in T84 cells. *Infect Immun* 71, 766-773.
- Dalmasso, G., Cottrez, F., Imbert, V., Lagadec, P., Peyron, J. F., Rampal, P., Czerucka, D., Groux, H., Foussat, A., and Brun, V. (2006a). *Saccharomyces boulardii* inhibits inflammatory bowel disease by trapping T cells in mesenteric lymph nodes. *Gastroenterology* 131, 1812-1825.
- Dalmasso, G., Loubat, A., Dahan, S., Calle, G., Rampal, P., and Czerucka, D. (2006b). *Saccharomyces boulardii*

- prevents TNF- $\alpha$ -induced apoptosis in EHEC-infected T84 cells. *Res Microbiol* 157, 456-465.
- de Llanos, R., Querol, A., Peman, J., Gobernado, M., and Fernandez-Espinar, M. T. (2006). Food and probiotic strains from the *Saccharomyces cerevisiae* species as a possible origin of human systemic infections. *Int J Food Microbiol* 110, 286-290.
- Dijkstra, G., Moshage, H., van Dullemen, H. M., de Jager-Krikken, A., Tiebosch, A. T., Kleibeuker, J. H., Jansen, P. L., and van Goor, H. (1998). Expression of nitric oxide synthases and formation of nitrotyrosine and reactive oxygen species in inflammatory bowel disease. *J Pathol* 186, 416-421.
- Dubuquoy, L., Jansson, E. A., Deeb, S., Rakotobe, S., Karoui, M., Colombel, J. F., Auwerx, J., Pettersson, S., and Desreumaux, P. (2003). Impaired expression of peroxisome proliferator-activated receptor gamma in ulcerative colitis. *Gastroenterology* 124, 1265-1276.
- Edwards-Ingram, L., Gitsham, P., Burton, N., Warhurst, G., Clarke, I., Hoyle, D., Oliver, S. G., and Stateva, L. (2007). Genotypic and physiological characterization of *Saccharomyces boulardii*, the probiotic strain of *Saccharomyces cerevisiae*. *Appl Environ Microbiol* 73, 2458-2467.
- Elmer, G. W., Surawicz, C. M., and McFarland, L. V. (1996). Biotherapeutic agents. A neglected modality for the treatment and prevention of selected intestinal and vaginal infections. *Jama* 275, 870-876.
- Escher, P., Braissant, O., Basu-Modak, S., Michalik, L., Wahli, W., and Desvergne, B. (2001). Rat PPARs: quantitative analysis in adult rat tissues and regulation in fasting and refeeding. *Endocrinology* 142, 4195-4202.
- FAO/WHO (2001). Evaluation of Health and Nutritional Properties of Probiotics in Food Including Powder Milk with Live Lactic Acid Bacteria (Cordoba, Argentina: Food and Agriculture organization of the United Nations and World Health Organization).
- Fietto, J. L., Araujo, R. S., Valadao, F. N., Fietto, L. G., Brandao, R. L., Neves, M. J., Gomes, F. C., Nicoli, J. R., and Castro, I. M. (2004). Molecular and physiological comparisons between *Saccharomyces cerevisiae* and *Saccharomyces boulardii*. *Can J Microbiol* 50, 615-621.
- Fuller, R. (1989). Probiotics in man and animals. *J Appl Bacteriol* 66, 365-378.
- Genolet, R., Wahli, W., and Michalik, L. (2004). PPARs as drug targets to modulate inflammatory responses? *Curr Drug Targets Inflamm Allergy* 3, 361-375.
- Girard, P., Pansart, Y., and Gillardin, J. M. (2005). Inducible nitric oxide synthase involvement in the mechanism of action of *Saccharomyces boulardii* in castor oil-induced diarrhoea in rats. *Nitric Oxide* 13, 163-169.
- Guandalini, S. (2000). Acute diarrhea. In *Pediatric Gastrointestinal Disease* (Walker WA, Durie PR, Hamilton JR, eds), B. D. Inc, ed., pp. 28-38.
- Guarner, F., and Malagelada, J. R. (2003). Gut flora in health and disease. *Lancet* 361, 512-519.
- Guslandi, M., Giollo, P., and Testoni, P. A. (2003). A pilot trial of *Saccharomyces boulardii* in ulcerative colitis. *Eur J Gastroenterol Hepatol* 15, 697-698.
- Guslandi, M., Mezzi, G., Sorghi, M., and Testoni, P. A. (2000). *Saccharomyces boulardii* in maintenance treatment of Crohn's disease. *Dig Dis Sci* 45, 1462-1464.
- Hennequin, C., Kauffmann-Lacroix, C., Jobert, A., Viard, J. P., Ricour, C., Jacquemin, J. L., and Berche, P. (2000). Possible role of catheters in *Saccharomyces boulardii* fungemia. *Eur J Clin Microbiol Infect Dis* 19, 16-20.
- Hennequin, C., Thierry, A., Richard, G. F., Lecoindre, G., Nguyen, H. V., Gaillardin, C., and Dujon, B. (2001). Microsatellite typing as a new tool for identification of *Saccharomyces cerevisiae* strains. *J Clin Microbiol* 39, 551-559.
- Herbrecht, R., and Nivoix, Y. (2005). *Saccharomyces cerevisiae* fungemia: an adverse effect of *Saccharomyces boulardii* probiotic administration. *Clin Infect Dis* 40, 1635-1637.
- Higgins, L. M., Frankel, G., Douce, G., Dougan, G., and MacDonald, T. T. (1999). *Citrobacter rodentium* infection in mice elicits a mucosal Th1 cytokine response and lesions similar to those in murine inflammatory bowel disease. *Infect Immun* 67, 3031-3039.
- Hogenauer, C., Hammer, H. F., Krejs, G. J., and Reisinger, E. C. (1998). Mechanisms and management of antibiotic-associated diarrhea. *Clin Infect Dis* 27, 702-710.
- Htwe, K., Yee, K. S., Tin, M., and Vandenplas, Y. (2008). Effect of *Saccharomyces boulardii* in the Treatment of Acute Watery Diarrhea in Myanmar Children: A Randomized Controlled Study. *Am J Trop Med Hyg* 78, 214-216.
- Jahn, H. U., Ullrich, R., Schneider, T., Liehr, R. M., Schieferdecker, H. L., Holst, H., and Zeitz, M. (1996). Immunological and trophical effects of *Saccharomyces boulardii* on the small intestine in healthy human volunteers. *Digestion* 57, 95-104.
- Jawahara, S., and Poulain, D. (2007). *Saccharomyces boulardii* decreases inflammation and intestinal colonization by *Candida albicans* in a mouse model of chemically-induced colitis. *Med Mycol*, 1-10.
- Kaper, J. B., Fasano, A., and Trucksis, M. (1994). Toxins of *Vibrio cholerae*. In *Vibrio cholerae* and cholera: molecular to global perspectives, I. K. Wachsmuth, P. A. Blake, and O. Olsvik, eds. (Washington, D.C.: American Society for Microbiology), pp. 145-176.
- Katz, D. A., Lynch, M. E., and Littenberg, B. (1996). Clinical prediction rules to optimize cytotoxin testing for *Clostridium difficile* in hospitalized patients with diarrhea. *Am J Med* 100, 487-495.
- Kelly, C. P., Pothoulakis, C., and LaMont, J. T. (1994). *Clostridium difficile* colitis. *N Engl J Med* 330, 257-262.
- Klein, S. M., Elmer, G. W., McFarland, L. V., Surawicz, C. M., and Levy, R. H. (1993). Recovery and elimination of the biotherapeutic agent, *Saccharomyces boulardii*, in healthy human volunteers. *Pharm Res* 10, 1615-1619.
- Kollaritsch, H., Holst, H., Grobara, P., and Wiedermann, G. (1993). [Prevention of traveler's diarrhea with *Saccharomyces boulardii*. Results of a placebo controlled double-blind study]. *Fortschr Med* 111, 152-156.
- Kotler, D. P., Gaetz, H. P., Lange, M., Klein, E. B., and Holt, P. R. (1984). Enteropathy associated with the acquired immunodeficiency syndrome. *Ann Intern Med* 101, 421-428.
- Kotowska, M., Albrecht, P., and Szajewska, H. (2005). *Saccharomyces boulardii* in the prevention of antibiotic-associated diarrhoea in children: a randomized double-blind placebo-controlled trial. *Aliment Pharmacol Ther* 21, 583-590.
- Lee, S. K., Kim, H. J., Chi, S. G., Jang, J. Y., Nam, K. D.,




- Kim, N. H., Joo, K. R., Dong, S. H., Kim, B. H., Chang, Y. W., et al. (2005). [*Saccharomyces boulardii* activates expression of peroxisome proliferator-activated receptor- $\gamma$  in HT-29 cells]. Korean J Gastroenterol 45, 328-334.
- Lherm, T., Monet, C., Nougier, B., Soulier, M., Larbi, D., Le Gall, C., Caen, D., and Malbrunot, C. (2002). Seven cases of fungemia with *Saccharomyces boulardii* in critically ill patients. Intensive Care Med 28, 797-801.
- Mansen, A., Guardiola-Diaz, H., Rafter, J., Branting, C., and Gustafsson, J. A. (1996). Expression of the peroxisome proliferator-activated receptor (PPAR) in the mouse colonic mucosa. Biochem Biophys Res Commun 222, 844-851.
- Maupas, J., Champemont, P., and Delforge, M. (1983). Treatment of irritable bowel syndrome with *Saccharomyces boulardii*: a double-blind, placebo-controlled study. Med Chir Dig 12, 77-79.
- McFarland, L. V. (1998). Epidemiology, risk factors and treatments for antibiotic-associated diarrhea. Dig Dis 16, 292-307.
- McFarland, L. V. (2006). Meta-analysis of probiotics for the prevention of antibiotic associated diarrhea and the treatment of *Clostridium difficile* disease. Am J Gastroenterol 101, 812-822.
- McFarland, L. V. (2007). Meta-analysis of probiotics for the prevention of traveler's diarrhea. Travel Med Infect Dis 5, 97-105.
- McFarland, L. V., Surawicz, C. M., Greenberg, R. N., Fekety, R., Elmer, G. W., Moyer, K. A., Melcher, S. A., Bowen, K. E., Cox, J. L., Noorani, Z., and et al. (1994). A randomized placebo-controlled trial of *Saccharomyces boulardii* in combination with standard antibiotics for *Clostridium difficile* disease. Jama 271, 1913-1918.
- Moncada, S., Palmer, R. M., and Higgs, E. A. (1991). Nitric oxide: physiology, pathophysiology, and pharmacology. Pharmacol Rev 43, 109-142.
- Mumy, K. L., Chen, X., Kelly, C. P., and McCormick, B. A. (2007). *Saccharomyces boulardii* interferes with *Shigella* pathogenesis by post-invasion signaling events. Am J Physiol Gastrointest Liver Physiol 294, G599-609.
- Ozkan, T. B., Sahin, E., Erdemir, G., and Budak, F. (2007). Effect of *Saccharomyces boulardii* in children with acute gastroenteritis and its relationship to the immune response. J Int Med Res 35, 201-212.
- Parkes, G. C., Brostoff, J., Whelan, K., and Sanderson, J. D. (2008). Gastrointestinal microbiota in irritable bowel syndrome: their role in its pathogenesis and treatment. Am J Gastroenterol 103, 1557-1567.
- Plein, K., and Hotz, J. (1993). Therapeutic effects of *Saccharomyces boulardii* on mild residual symptoms in a stable phase of Crohn's disease with special respect to chronic diarrhea--a pilot study. Z Gastroenterol 31, 129-134.
- Pothoulakis, C. (1996). Pathogenesis of *Clostridium difficile*-associated diarrhoea. Eur J Gastroenterol Hepatol 8, 1041-1047.
- Qamar, A., Aboudola, S., Warny, M., Michetti, P., Pothoulakis, C., LaMont, J. T., and Kelly, C. P. (2001). *Saccharomyces boulardii* stimulates intestinal immunoglobulin A immune response to *Clostridium difficile* toxin A in mice. Infect Immun 69, 2762-2765.
- Rodrigues, A. C., Cara, D. C., Fretez, S. H., Cunha, F. Q., Vieira, E. C., Nicoli, J. R., and Vieira, L. Q. (2000). *Saccharomyces boulardii* stimulates sIgA production and the phagocytic system of gnotobiotic mice. J Appl Microbiol 89, 404-414.
- Roy, C., Silverman, A., and Alagille, D. (1995). Diarrheal disorders, In Pediatric Clinical Gastroenterology (St Louis: Mosby), pp. 216-287.
- Saint-Marc, T., Rossello-Prats, L., and Touraine, J. L. (1991). [Efficacy of *Saccharomyces boulardii* in the treatment of diarrhea in AIDS]. Ann Med Interne (Paris) 142, 64-65.
- Sakaguchi, S., Sakaguchi, N., Asano, M., Itoh, M., and Toda, M. (1995). Immunologic self-tolerance maintained by activated T cells expressing IL-2 receptor  $\alpha$ -chains (CD25). Breakdown of a single mechanism of self-tolerance causes various autoimmune diseases. J Immunol 155, 1151-1164.
- Sakaguchi, S., Wing, K., and Miyara, M. (2007). Regulatory T cells - a brief history and perspective. Eur J Immunol 37 Suppl 1, S116-123.
- Sanders, J. W., and Tribble, D. R. (2001). Diarrhea in the returned traveler. Curr Gastroenterol Rep 3, 304-314.
- Schauer, D. B., and Falkow, S. (1993). Attaching and effacing locus of a *Citrobacter freundii* biotype that causes transmissible murine colonic hyperplasia. Infect Immun 61, 2486-2492.
- Schlotterer, M., Bernasconi, P., Lebreton, F., and al., e. (1987). Intérêt de *Saccharomyces Boulardii* dans la tolérance digestive de la nutrition entérale à débit continu chez le brûlé. Nutr Clin Metabol 1, 31-34.
- Schneider, S. M., Girard-Pipau, F., Filippi, J., Hebuterne, X., Moyse, D., Hinojosa, G. C., Pompei, A., and Rampal, P. (2005). Effects of *Saccharomyces boulardii* on fecal short-chain fatty acids and microflora in patients on long-term total enteral nutrition. World J Gastroenterol 11, 6165-6169.
- Schneider, S. M., Le Gall, P., Girard-Pipau, F., Piche, T., Pompei, A., Nano, J. L., Hebuterne, X., and Rampal, P. (2000). Total artificial nutrition is associated with major changes in the fecal flora. Eur J Nutr 39, 248-255.
- Sougioultzis, S., Simeonidis, S., Bhaskar, K. R., Chen, X., Anton, P. M., Keates, S., Pothoulakis, C., and Kelly, C. P. (2006). *Saccharomyces boulardii* produces a soluble anti-inflammatory factor that inhibits NF- $\kappa$ B-mediated IL-8 gene expression. Biochem Biophys Res Commun 343, 69-76.
- Strober, W., Fuss, I., and Mannon, P. (2007). The fundamental basis of inflammatory bowel disease. J Clin Invest 117, 514-521.
- Su, C. G., Wen, X., Bailey, S. T., Jiang, W., Rangwala, S. M., Keilbaugh, S. A., Flanagan, A., Murthy, S., Lazar, M. A., and Wu, G. D. (1999). A novel therapy for colitis utilizing PPAR- $\gamma$  ligands to inhibit the epithelial inflammatory response. J Clin Invest 104, 383-389.
- Surawicz, C. M., McFarland, L. V., Greenberg, R. N., Rubin, M., Fekety, R., Mulligan, M. E., Garcia, R. J., Brandmarker, S., Bowen, K., Borjal, D., and Elmer, G. W. (2000). The search for a better treatment for recurrent *Clostridium difficile* disease: use of high-dose vancomycin combined with *Saccharomyces boulardii*. Clin Infect Dis 31, 1012-1017.
- Szajewska, H., Ruszczynski, M., and Radzikowski, A. (2006). Probiotics in the prevention of antibiotic-associated diarrhea in children: a meta-analysis of randomized



- controlled trials. *J Pediatr* 149, 367-372.
- Szajewska, H., Skorka, A., and Dylag, M. (2007). Meta-analysis: *Saccharomyces boulardii* for treating acute diarrhoea in children. *Aliment Pharmacol Ther* 25, 257-264.
- Tasteyre, A., Barc, M. C., Karjalainen, T., Bourlioux, P., and Collignon, A. (2002). Inhibition of *in vitro* cell adherence of *Clostridium difficile* by *Saccharomyces boulardii*. *Microb Pathog* 32, 219-225.
- Tempe, J., Steidel, A., Bléhaut, H., and *al.*, e. (1983). Prévention par *Saccharomyces boulardii* des diarrhées de l'alimentation entérale à débit continu. *Sem Hop Paris* 59, 1409-1412.
- Torres, J., Jennische, E., Lange, S., and Lonnroth, I. (1990). Enterotoxins from *Clostridium difficile*; diarrhoeogenic potency and morphological effects in the rat intestine. *Gut* 31, 781-785.
- Triadafilopoulos, G., Pothoulakis, C., O'Brien, M. J., and LaMont, J. T. (1987). Differential effects of *Clostridium difficile* toxins A and B on rabbit ileum. *Gastroenterology* 93, 273-279.
- Triadafilopoulos, G., Pothoulakis, C., Weiss, R., Giampaolo, C., and Lamont, J. T. (1989). Comparative study of *Clostridium difficile* toxin A and cholera toxin in rabbit ileum. *Gastroenterology* 97, 1186-1192.
- Vidon, N., Huchet, B., and Rambaud, J. C. (1986). [Influence of *Saccharomyces boulardii* on jejunal secretion in rats induced by cholera toxin]. *Gastroenterol Clin Biol* 10, 13-16.
- Vollaard, E. J., and Clasener, H. A. (1994). Colonization resistance. *Antimicrob Agents Chemother* 38, 409-414.
- Whelan, K., Gibson, G. R., Judd, P. A., and Taylor, M. A. (2001). The role of probiotics and prebiotics in the management of diarrhoea associated with enteral tube feeding. *J Hum Nutr Diet* 14, 423-433.
- Wu, X., Vallance, B. A., Boyer, L., Bergstrom, K. S., Walker, J., Madsen, K., O'Kusky, J. R., Buchan, A. M., and Jacobson, K. (2008). *Saccharomyces boulardii* ameliorates *Citrobacter rodentium*-induced colitis through actions on bacterial virulence factors. *Am J Physiol Gastrointest Liver Physiol* 294, G295-306.
- Xavier, R. J., and Podolsky, D. K. (2007). Unravelling the pathogenesis of inflammatory bowel disease. *Nature* 448, 427-434.

### Further Reading

- Ljungh, A. and Wadström, T. (2009). *Lactobacillus Molecular Biology: From Genomics to Probiotics* (Norfolk: Caister Academic Press).  
<http://www.caister.com/probiotics>
- Tannock, G.W. (2005). *Probiotics and Prebiotics: Scientific Aspects* (Norfolk: Caister Academic Press).  
<http://www.caister.com/probiotics>
- Tannock, G.W. (2002). *Probiotics and Prebiotics: Where are We Going?* (Norfolk: Caister Academic Press).  
<http://www.caister.com/probiotics>



## ***Lactobacillus* Molecular Biology**

### **From Genomics to Probiotics**

**Publisher:** Caister Academic Press

**Edited by:** Åsa Ljungh and Torkel Wadström

**x + 206 pp., January 2009**




**ISBN 978-1-904455-41-7 \$310 / £150**

Includes phylogenetics, taxonomy, comparative genomics, functional genomics, intestinal microflora, surface proteins, stress responses, immune system, probiotics, anti-cancer potential, and much more. Essential reading for all scientists involved in lactic acid bacteria or probiotic research and a recommended book for all microbiology laboratories.




# Further Reading

**Caister Academic Press** is a leading academic publisher of advanced texts in microbiology, molecular biology and medical research. Full details of all our publications at [caister.com](http://www.caister.com)

- **MALDI-TOF Mass Spectrometry in Microbiology**  
Edited by: M Kostrzewa, S Schubert (2016)  
[www.caister.com/malдитof](http://www.caister.com/malдитof)
- ***Aspergillus* and *Penicillium* in the Post-genomic Era**  
Edited by: RP Vries, IB Gelber, MR Andersen (2016)  
[www.caister.com/aspergillus2](http://www.caister.com/aspergillus2)
- **The Bacteriocins: Current Knowledge and Future Prospects**  
Edited by: RL Dorit, SM Roy, MA Riley (2016)  
[www.caister.com/bacteriocins](http://www.caister.com/bacteriocins)
- **Omics in Plant Disease Resistance**  
Edited by: V Bhadauria (2016)  
[www.caister.com/opdr](http://www.caister.com/opdr)
- **Acidophiles: Life in Extremely Acidic Environments**  
Edited by: R Quatrini, DB Johnson (2016)  
[www.caister.com/acidophiles](http://www.caister.com/acidophiles)
- **Climate Change and Microbial Ecology: Current Research and Future Trends**  
Edited by: J Marxsen (2016)  
[www.caister.com/climate](http://www.caister.com/climate)
- **Biofilms in Bioremediation: Current Research and Emerging Technologies**  
Edited by: G Lear (2016)  
[www.caister.com/biorem](http://www.caister.com/biorem)
- **Microalgae: Current Research and Applications**  
Edited by: MN Tsaloglou (2016)  
[www.caister.com/microalgae](http://www.caister.com/microalgae)
- **Gas Plasma Sterilization in Microbiology: Theory, Applications, Pitfalls and New Perspectives**  
Edited by: H Shintani, A Sakudo (2016)  
[www.caister.com/gasplasma](http://www.caister.com/gasplasma)
- **Virus Evolution: Current Research and Future Directions**  
Edited by: SC Weaver, M Denison, M Roossinck, et al. (2016)  
[www.caister.com/virusevol](http://www.caister.com/virusevol)
- **Arboviruses: Molecular Biology, Evolution and Control**  
Edited by: N Vasilakis, DJ Gubler (2016)  
[www.caister.com/arbo](http://www.caister.com/arbo)
- ***Shigella*: Molecular and Cellular Biology**  
Edited by: WD Picking, WL Picking (2016)  
[www.caister.com/shigella](http://www.caister.com/shigella)
- **Aquatic Biofilms: Ecology, Water Quality and Wastewater Treatment**  
Edited by: AM Romani, H Guasch, MD Balaguer (2016)  
[www.caister.com/aquaticbiofilms](http://www.caister.com/aquaticbiofilms)
- **Alphaviruses: Current Biology**  
Edited by: S Mahalingam, L Herrero, B Herring (2016)  
[www.caister.com/alpha](http://www.caister.com/alpha)
- **Thermophilic Microorganisms**  
Edited by: F Li (2015)  
[www.caister.com/thermophile](http://www.caister.com/thermophile)



- **Flow Cytometry in Microbiology: Technology and Applications**  
Edited by: MG Wilkinson (2015)  
[www.caister.com/flow](http://www.caister.com/flow)
- **Probiotics and Prebiotics: Current Research and Future Trends**  
Edited by: K Venema, AP Carmo (2015)  
[www.caister.com/probiotics](http://www.caister.com/probiotics)
- **Epigenetics: Current Research and Emerging Trends**  
Edited by: BP Chadwick (2015)  
[www.caister.com/epigenetics2015](http://www.caister.com/epigenetics2015)
- ***Corynebacterium glutamicum*: From Systems Biology to Biotechnological Applications**  
Edited by: A Burkovski (2015)  
[www.caister.com/cory2](http://www.caister.com/cory2)
- **Advanced Vaccine Research Methods for the Decade of Vaccines**  
Edited by: F Bagnoli, R Rappuoli (2015)  
[www.caister.com/vaccines](http://www.caister.com/vaccines)
- **Antifungals: From Genomics to Resistance and the Development of Novel Agents**  
Edited by: AT Coste, P Vandeputte (2015)  
[www.caister.com/antifungals](http://www.caister.com/antifungals)
- **Bacteria-Plant Interactions: Advanced Research and Future Trends**  
Edited by: J Murillo, BA Vinatzer, RW Jackson, et al. (2015)  
[www.caister.com/bacteria-plant](http://www.caister.com/bacteria-plant)
- ***Aeromonas***  
Edited by: J Graf (2015)  
[www.caister.com/aeromonas](http://www.caister.com/aeromonas)
- **Antibiotics: Current Innovations and Future Trends**  
Edited by: S Sánchez, AL Demain (2015)  
[www.caister.com/antibiotics](http://www.caister.com/antibiotics)
- ***Leishmania*: Current Biology and Control**  
Edited by: S Adak, R Datta (2015)  
[www.caister.com/leish2](http://www.caister.com/leish2)
- ***Acanthamoeba*: Biology and Pathogenesis (2nd edition)**  
Author: NA Khan (2015)  
[www.caister.com/acanthamoeba2](http://www.caister.com/acanthamoeba2)
- **Microarrays: Current Technology, Innovations and Applications**  
Edited by: Z He (2014)  
[www.caister.com/microarrays2](http://www.caister.com/microarrays2)
- **Metagenomics of the Microbial Nitrogen Cycle: Theory, Methods and Applications**  
Edited by: D Marco (2014)  
[www.caister.com/n2](http://www.caister.com/n2)



Order from [caister.com/order](http://caister.com/order)